
EFDA–JET–CP(01)02-04

V.Philipps, R.A.Pitts, G.Matthews, M.De Baar, J.Cordey, T.Eich,
M.Fenstermacher, W.Fundamenski, A.Herrmann, D.Hillis, A.Huber,
T.Loarer, A.Loarte, P.Lomas, D.C.McDonald, A.Pospieszczyk, J.Rapp,
V.Riccardo, E.Righi, G.Saibene, M.Stamp, J.Strachan, K.D.Zastrow
and JET EFDA Contributors

Recent Results from Helium Plasma Operation in JET

Recent Results from Helium Plasma Operation in JET

V.Philipps¹, R.A.Pitts², G.Matthews³, M.De Baar⁹, J.Cordey³, T.Eich¹,
M.Fenstermacher⁴, W.Fundamenski³, A.Herrmann⁶, D.Hillis⁵, A.Huber¹,
T.Loarer⁸, A.Loarte⁷, P.Lomas³, D.C.McDonald³, A.Pospieszczyk¹, J.Rapp¹,
V.Riccardo, E.Righi⁷, G.Saibene⁷, M.Stamp³, J.Strachan¹⁰, K.D.Zastrow³
and JET EFDA Contributors*

¹*Institut für Plasmaphysik, Association EURATOM, Forschungszentrum Jülich, Germany,*

²*CRPP/EPFL, Association EURATOM-Confederation Suisse, CH-1015 Lausanne, Switzerland,*

³*Euratom UKAEA Fusion Association, Culham Science Centre, Abingdon, Oxon OX14 3DB, UK,*

⁴*DIII-D National Fusion facility, San Diego, USA,*

⁵*Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA,*

⁶*Max-Planck-Institut für Plasmaphysik, Garching, Germany,*

⁷*EFDA-CSU, D-85748 Garching, Germany,*

⁸*Association EURATOM-CEA, DRFC, CEA Cadarache, France. Sweden*

⁹*FOM-Rijnhuizen, Association EURATOM-FOM, Nieuwegein,*

¹⁰*PPPL, Princeton University, USA*

**See appendix of the paper by J.Pamela "Overview of recent JET results",
Proceedings of the IAEA conference on Fusion Energy, Sorrento 2000*

Preprint of Paper to be submitted for publication in Proceedings of the
EPS Conference,
(Madeira, Portugal 18-22 June 2001)

“This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

“Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

INTRODUCTION

To assess the suitability for ITER of operation during the low activation phase in ELMing H-mode helium plasmas and to investigate in detail areas in which the underlying physics differs from operation with hydrogenic isotopes as the main plasma species, pure H_e discharges have been explored in JET during a three week period at the end of first experimental campaign in 2001. As many as possible of the principal operating regimes in D (e.g. L and H-mode and OS plasmas) have been explored with the aim of improving understanding in confinement, the L-H transition, ELMs and edge physics. With regard to the latter, particular topics of interest include atomic physics and impurity production and their influence on divertor radiation, detachment and density limits.

1. EXPERIMENTAL AND OPERATIONAL CONDITIONS

For much of the H_e campaign both JET beam boxes were converted for pure helium injection using Ar frosting on the NI-cryopumps. During D to H_e changeover using a fixed magnetic equilibrium and a reproducible ICRH power ramp to investigate L-H transition thresholds, H_e/D ratios $>85\%$ were obtained after only 10 discharges. Changing the configuration increased the D plasma content even after relatively long term operation in H_e , demonstrating that a significant D reservoir exists in surfaces not previously exposed to direct H_e ion fluxes. Although the lack of active H_e pumping in most discharges often yielded uncontrolled increases in core and edge densities with subsequent influence on, for example, confinement, ELMs and access to OS scenarios, a few discharges with divertor cryopump Ar frosting were performed in which steady state H-Mode conditions were achieved.

2. CONFINEMENT AND L-H TRANSITION

For the first time a set of quasi steady state ELMy-H mode shots have been achieved with H_e -NBI injection up to 12MW in discharges with I_p , BT and the ratio H_e/D in the range 1-2MA, 1-2T and 70-90% respectively. Without active H_e cryopumping, the density rise leads in general to a type I-III ELM transition as shown in Fig1. Although the effect of D in H_e plasmas is still to be assessed, preliminary analysis shows that the L-H threshold is about 50% higher with respect to that in D, with a similar dependence on I_p and BT. The energy confinement decreases by $\sim 15-20\%$ (see Fig. 2) compared with equivalent D plasmas, which would result in a confinement dependence of $Z^{-0.55}$ assuming a positive mass dependence ($\sim M^{0.2}$) as measured in previous JET isotope measurements (1).

3. IMPURITY PRODUCTION AND DIVERTOR PHYSICS

In comparison with similar D plasmas, the carbon impurity release in H_e discharges changes significantly for all conditions studied. This highlights the importance of chemical erosion and Charge eXchange (CX) neutral fluxes, both of which change considerably during H_e operation. Fig.3 compares the CIII light emission integrated over the inner and outer divertor legs and that along a midplane horizontal chord from two similar L-mode density limit shots (a) and two 11MW H-mode discharges at the same density (b) in H_e and D. There is a large (by factors 3-10) decrease

in the inner divertor carbon source, an observation common to all plasma conditions studied. In the outer divertor the carbon release changes little at lower plasma densities but, in contrast to D operation, decreases steadily with increasing density. A horizontal spectroscopic view recorded similar CIII light emission during the limiter phases, but a dramatic reduction in the carbon light during the diverted phase in H_e. Other C-ionisation states behave similarly in the different plasma scenarios. The data demonstrate that in the inner divertor chemical erosion largely dominates the carbon source in deuterium plasmas. By contrast, at the outer divertor physical sputtering dominates at low density, with chemical erosion becoming increasingly important as density increases. In the main chamber the reduction of the carbon source in helium plasmas is mainly due to reduced CX fluxes.

In L-mode H_e plasmas, the reduced carbon influx results in a considerable reduction in main plasma carbon content. This is clearly illustrated in Fig. 3a by the strong decrease in the incremental Z_{eff} compared with a similar D plasma and also by the carbon concentrations from CX spectroscopy which are below the detection limit at higher plasma densities in H_e. In H-mode, the carbon source reduction is similar but this is not manifest in a reduced Z_{eff} (Fig 3b). Such different behaviour is surprising since it implies an increasing influence of the outer target carbon source on the main plasma C content during H-mode conditions.

Helium operation and the consequent reduced carbon influxes lead to L-mode density limits about a factor of two higher than those found for identical D plasmas (Fig. 3a). Increasing the density in a deuterium discharge leads to detachment of the inner divertor leg with subsequent strong X-point radiation and finally to the development of an inner wall MARFE resulting in a density limit disruption (2). In helium, the L-mode density limit is characterised by an almost poloidally symmetric radiative collapse ($P_{\text{rad}} = P_{\text{in}}$), which results also in a different disruption behaviour(3).

The longer H_e ionisation mean-free-path allows neutrals to penetrate the X-point before the low divertor temperatures required for recombination to occur can be reached, leading to leading to substantially modified detachment behaviour at high density in comparison with D discharges. Radiation otherwise originating from low charge states of carbon in D plasmas is presently thought to originate from H_eII line emission during helium operation. Interestingly, at a given density and within a certain power range, periodic oscillations of the H_eI light emission in the inner divertor have been observed in L-mode. This phenomena is probably associated with the radiation behaviour of H_e leading to T_e oscillations when a critical temperature and density is reached.

Power deposition on the divertor targets has been investigated by IR thermography, thermocouples and Langmuir probes. In general, the inner/outer target asymmetry in this deposition is similar in H_e to that in D (Fig 4). Preliminary analysis shows that the deposition profiles are 30-80% larger than in deuterium. Investigation of the ELM behaviour is of particular interest for the validation of ELM models since collisionality in H_e is four times that in D (at the same T_e). The data indicate that the ELM rise time (~500-800s) is larger than in comparable D discharges, although studies of ELMs in D show that the ELM rise time can vary significantly (4).

SUMMARY

Helium plasmas in L- and H-mode (1-2MA, 1-2T) with H_e NBI up to 12MW have been studied in JET. Type I and III ELMy H-modes have been routinely obtained. Energy confinement is observed to decrease by about 20% in H_e compared with similar D plasmas and the L-H transition power is about 50% higher in H_e. Achievement of H-mode in D plasmas is therefore easier than in H_e and this is an important conclusion for the non-active phase of ITER. Carbon impurity release at the inner divertor is significantly decreased in H_e, showing that chemical erosion in D plasmas determines more or less completely the carbon production there. At the outer target, physical sputtering is an important C source in helium, particularly during H-mode. Reduced CX fluxes in H_e lead to a strong decrease of carbon released from main chamber wall in diverted discharges. The H_e L-mode density limit is about twice that in D with the limit being determined by overall radiation collapse. In general, although relevant ELMy H-modes can be obtained in helium, helium operation would not provide an adequate test of critical divertor physics issues for ITER which must be a key objective for any low activation phase.

REFERENCES

- [1]. Nuclear Fusion, Vol. 39, No. 9 (1999) 1133
- [2]. J. Rapp et al this conference
- [3]. V.Riccardo, P.Andrew , K.H.Finken to be published
- [4]. T.Eich et al , this conference




Figure 1: Typical behaviour of density and ELM behaviour in a H_e ELMy H-mode without active He pumping




Figure 2: Confinement based on $H_{98}(y,2)$ without mass dependence for D and H_e pulses in the range of 1-2 MA





Figure 3(a): Evolution of the density, CIII light in the outer, inner divertor and midplane and Z_{eff} for L-mode density limit shots in H_e and D (low wall clearance). (b) Evolution of the CIII light emission and Z for two Elmy H-mode shots in H_e and D at 11 MW input power





Figure 4(a): Distribution of total energy over the divertor target tiles derived from thermocouple data for two similar L-mode shots in H_e and D. (b) Power decay lengths for the inner and outer divertor transformed to the horizontal midplane for L-mode shots in D and H_e and different power levels. Data are derived from thermocouple and Langmuir probe data