
EFDA–JET–CP(02)07/05

X. Garbet, Y. Baranov, G. Bateman, S. Benkadda, P. Beyer, R. Budny, F. Crisanti,
B. Esposito, C. Figarella, C. Fourment, P. Ghendrih, F. Imbeaux, E. Joffrin,
J. Kinsey, A. Kritz, X. Litaudon, P. Maget, P. Mantica, D. Moreau, Y. Sarazin,
A. Pankin, V. Parail, A. Peeters, T. Tala, G. Tardini, A. Thyagaraja,
I. Voitsekhovitch, J. Weiland and R. Wolf

Micro-Stability and Transport Modelling of Internal Transport Barriers on JET

Micro-Stability and Transport Modelling of Internal Transport Barriers on JET

X. Garbet¹, Y. Baranov², G. Bateman³, S. Benkadda⁴, P. Beyer⁴, R. Budny⁵,
F. Crisanti⁶, B. Esposito⁶, C. Figarella⁴, C. Fourment¹, P. Ghendrih¹, F. Imbeaux¹,
E. Joffrin¹, J. Kinsey³, A. Kritz³, X. Litaudon¹, P. Maget¹, P. Mantica⁷, D. Moreau¹,
Y. Sarazin¹, A. Pankin³, V. Parail², A. Peeters¹⁰, T. Tala⁹, G. Tardini¹⁰
A. Thyagaraja², I. Voitsekhovitch⁴, J. Weiland⁸, R. Wolf¹¹
and contributors to the EFDA-JET workprogramme*

¹Association Euratom-CEA, CEA Cadarache, 13108 St Paul-Lez-Durance, France.

²EURATOM/UKAEA, Culham Science Centre, Abingdon OX14 3DB, United Kingdom.

³Lehigh University Physics Department, 16 Memorial Drive East, Bethlehem, PA 18015, USA.

⁴LPIIM, Centre Universitaire de Saint-Jerôme, 13397 Marseille cedex 20, France.

⁵PPPL, Princeton University, P.O. Box 451, Princeton, NJ 08543, USA.

⁶Associazione EURATOM-ENEA sulla Fusione, Via Enrico Fermi 27, 00044 Frascati, Italy.

⁷Istituto di Fisica del Plasma CNR-EURATOM, via Cozzi 53, 20125 Milano, Italy.

⁸Chalmers University of Technology and Euratom-VR Association, S-41296 Göteborg, Sweden.

⁹Association EURATOM-TEKES, VTT CTIP, FIN-02044 VTT, Finland.

¹⁰MPI für Plasmaphysik, EURATOM-Assoziation, D-8046 Garching bei München, Germany

¹¹Institut für Plasmaphysik, Association EURATOM / FZJ, D-52425 Jülich

* See annex of J. Pamela et al, "Overview of Recent JET Results and Future Perspectives",
Fusion Energy 2000 (Proc. 18th Int. Conf. Sorrento, 2000), IAEA, Vienna (2001).

“This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

“Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

ABSTRACT.

The physics of ITB formation in JET has been investigated using micro-stability analysis, profile modelling and turbulence simulations. The calculation of linear growth rates show that the magnetic shear plays a crucial role in the formation of the ITB. The Shafranov shift, ratio of the ion to electron temperature, and impurity content further improve the stability. This picture is consistent with profile modelling and global fluid simulations of electrostatic drift waves. Turbulence simulations also show that rational q values may play a special role in triggering an ITB. The same physics also explains how double internal barriers can be formed.

1. INTRODUCTION

Internal Transport Barriers (ITB's) in tokamak plasmas are considered as a promising way to achieve steady-state plasmas with good confinement properties in a fusion reactor. A crucial question is whether it will be possible to produce an ITB in a next step device with a reasonable amount of power. Once a barrier is triggered, a self-amplifying process takes place, where increasing gradients produce $E \times B$ velocity shear and Shafranov shift large enough to further decrease the turbulent transport. This paper is however focused on the question of barrier initiation. Many experimental results in JET point towards the safety factor profile as a key ingredient. In particular the power threshold is clearly lower when the magnetic shear is reversed. However other mechanisms like Shafranov shift stabilisation, impurity content or density peaking may play a role. One aim of this paper is to apply and compare various models and techniques on a common set of JET plasmas. Micro-stability analysis, profile modelling and turbulence simulations are used to this purpose. This paper also tackles a challenging class of transport barriers that are sensitive to low order rational surfaces. Their role has been recently confirmed in JET reversed shear plasmas, thanks to the observation of Alfvén cascades in coincidence with barrier formation. In particular strong barriers are often triggered when q_{\min} crosses 2 or 3. Surprisingly when q_{\min} further decreases with time and falls below $q=2$, the barrier sometimes splits. Two internal barriers then coexist and are tied to the $q=2$ magnetic surfaces. Existing models are confronted to this puzzling behaviour.

2. BRIEF DESCRIPTION OF JET ITB'S.

The physics that is usually invoked for explaining the triggering and self-sustainment of an ITB is a mixture of turbulence suppression via $E \times B$ velocity shear and linear stabilisation of drift waves. The magnetic shear is often considered as the main reason for improved stability. Two mechanisms have been identified: a decrease of the interchange drive [1], which is more prominent at negative shear, and a rarefaction of resonant surfaces that occurs at zero shear [2]. The reduction in turbulent transport comes from a decrease of the drive and/or smaller correlation lengths. Other parameters likely play some role such as the Shafranov shift (also called α effect), density gradient, impurity content and ratio of the ion to electron temperature. A common way to assess this stabilisation is to compare the $E \times B$ shear rate γ_E to a maximum linear growth rate γ_{lin} [3,4,5].

Two operational criteria can be built on the basis of this simple rule. Assuming $T_e \approx T_i$, the linear growth rate γ_{lin} of ITG/TEM modes is of the form c_s/L_{Te} up to a function of plasma parameters (c_s is the sound speed and L_{Te} the electron temperature gradient length). The diamagnetic part of the rotational shear rate reads $\gamma_E \approx \rho_S c_s / L_{Te}^2$. Including the contribution of the toroidal velocity in an effective linear growth rate, the criterion for a first order transition is $\rho_T^* = \rho_S / L_{Te} > \rho_{\text{Tcrit}}^*$. In principle ρ_{Tcrit}^* depends on the magnetic shear and the Mach number. In practice an analysis of the JET database shows that this criterion works well with a constant $\rho_{\text{Tcrit}}^* = 0.014$ [7]. Another criterion corresponds to a “loss of stiffness”. Stiffness means here that the temperature gradient length (for ions or electrons) is close to a threshold value $R/L_T = R/L_{T\text{crit}}$. This hypothesis is still under investigation at JET. Ion Cyclotron modulation experiments with mode conversion in L mode show the existence of a threshold for electrons [8]. For ions evidence has been obtained from steady-state profiles in L and H modes [9]. A natural definition of an ITB then corresponds to a region where the threshold is well above the L mode value. This leads to a criterion of the form $R/L_T > R/L_{T\text{crit}}$ for the ITB formation. This rule can be written as a condition on the ratio of core to edge temperature. A large class of ion ITB’s was found to satisfy this criterion using a critical value $R/L_{T\text{crit}} \approx 6$ [9].

3. MICRO-STABILITY ANALYSIS OF JET PLASMAS WITH AN ITB.

Several fluid and kinetic stability codes have been used to calculate the growth rates of Ion Temperature Gradient (ITG) modes and Trapped Electron Modes (TEM) [10-17]. The various techniques used to calculate the linear growth rates and $E \times B$ velocity shear are summarised in Table I. All groups have used the Hahn-Burrell definition of the $E \times B$ shear rate. However the calculation procedure was different (see Table I), thus leading to substantial differences (Fig.1). These models have been compared on the same JET plasma Pulse No: 51976. This pulse is a transient ITB with high performance that was analysed in detail by Challis et al. [22].

Name	Growth rate	E_r calculation
Weiland [10]	Fluid ITG (Weiland [14])	NCLASS [19] Flux coordinate
GS2 [11]	gyrokinetic flux tube ITG/TEM (GS2) [16]	NCLASS [19] Local
Rogister [12]	Rogister model [15]	Kim model [20] Local
Kine0 [13]	variational gyrokinetic ITG/TEM (KINEZERO) [18]	Kim model [20] Local

Table I: List of models used to analyse JET transport barriers.

The q profile is reversed early in the discharge with the help of current pre-forming phase with Lower Hybrid Current Drive (LHCD). A barrier forms early in the plasma at $t \approx 1.5$ s, after LHCD is applied,

and is strongly amplified at $t \approx 6$ s (see Fig.1). The q profile shown in Fig.1 is from a TRANSP run [11]. The result of the stability analysis at $t=6$ s, before the barrier strengthens, is shown on Fig. 1. Note that this time slice takes place where a barrier already exists so that the velocity shear rate is large. Three models do predict stabilisation, whereas the Weiland model predicts growth rates that are too large to be overcome by the velocity shear rate. However this model uses a ballooning representation that is not valid in the vicinity of $q=q_{\min}$. Using the Rogister model [15] instead gives a better agreement. Explaining the barrier onset at $t=1.5$ s is much more difficult. A transition due to the $E \times B$ velocity shear or a stabilisation alone is not possible. Indeed the shear rate is too low ($\approx 10^4 \text{ s}^{-1}$) compared to a typical value of γ_{lin} , unless a burst of localised rotational shear occurs, as observed in TFTR (this possibility is analysed in §6). Thus a decrease of the growth rate has to be invoked to explain this transition. In practice, most models rely essentially on the magnetic shear to trigger the barrier via a decrease of the linear growth rate. This effect is less marked when using the GS2 code, which predicts a transition at $t \approx 4$ s [11]. In the latter case, the stabilisation is due to the combined contributions of the negative magnetic shear, Shafranov shift and impurity content. No obvious difference is seen between negative and zero magnetic shear. The Rogister model favours low magnetic shear, confirmed by a recent analysis of the JET database [23], whereas the GS2 (flux tube) code seems to be more sensitive to negative shear.

4. PROFILE MODELLING OF JET ITB'S.

JET ITB plasmas have been modelled using several available transport models: Mixed Bohm-gyroBohm (B/gB) [24,25], Multi-Mode (MMM) [26], and Weiland [14,27] models. Bohm/gyroBohm models have been implemented in the JETTO and CRONOS codes. The main differences between the two codes come from the LHCD modules (FRFC in JETTO and Delphine in CRONOS). Moreover the stabilisation by magnetic shear and $E \times B$ velocity is implemented in different ways. Namely the JETTO codes enforces a global decrease of the diffusivity in the region where $\gamma_E > 0.68\gamma_{\text{ITG}}(s-0.14)$ [24], where γ_{ITG} is approximated by v_{Ti}/R (v_{Ti} is the ion thermal velocity). The CRONOS local uses a smoother and local reduction of the diffusivity $1/[1 + \exp(20(0.05 + \gamma_E/\gamma_{\text{ITG}} - s))]$, with a growth rate γ_{ITG} given by Newman et al. [17]. This exercise was carried out for the Pulses Nos:51976 (see §2) and the quasi-steady state ITB Pulse No: 53521 with LHCD throughout the pulse ([28]), with similar results. The whole pulse history was simulated. A comparison is shown in Fig.2 in the steady-state phase.

The best agreement is reached for the models that are very sensitive to the magnetic shear. Interestingly the two simulations using the Mixed Bohm/gyroBohm model do show some differences. This is due to the different current drive modules and the differences in the nature of the transition (local or global). This sensitivity to the current profile is not surprising since the onset of the barrier is mainly due to the magnetic shear, whereas the velocity shear rate is small at the transition. This result is in line with the findings of the stability analysis (§2). Later on in the pulse, the velocity shear rate becomes increasingly important for maintaining the barrier and moving its location

outward. The Multi-Mode model also follows the same trend (in particular the magnetic shear appears explicitly via the Hamaguchi-Horton definition of the rotational shear). Weiland model predicts ITB formation but the density gradient seems to be the key ingredient in this case [27]. Thus, although many results point in the direction of the magnetic shear as the main responsible of the transition to an ITB, other mechanisms cannot be excluded.

5. TURBULENCE SIMULATIONS OF JET ITB'S.

Global fluid simulations of electrostatic ITG/TEM modes (TRB code,[29,30]) have been run for several JET plasmas. All simulations show the importance of the magnetic shear for the onset of the barrier. However different mechanisms are involved for ions and electrons. Although high wave number ITG modes are stabilised by negative shear, the main reason for the onset of an ion barrier is the formation of a gap in the density of rational surfaces at low wave numbers close to the minimum of safety factor q_{\min} (see Fig.3). A barrier appears when this gap is larger than a turbulence correlation length (of the order of a centimetre in JET). Once an ion barrier is produced, its position and width are controlled by rotational shear. Electrons are sensitive to both negative and zero magnetic shear. Obviously TEMs are also affected by a gap in the resonant surfaces, even if they do not need an overlap with adjacent resonant surfaces to be unstable. TEMs are also affected by a negative magnetic shear because of the reversal of the trapped electron curvature drift that decreases the instability drive. Full stabilisation of TEM modes is expected for $s < -3/8$ in this model.

Simulations of an actual ITB in JET indicate that all mechanisms are involved, depending on the q profile. The example of the Pulse No: 53521 is shown in Fig.3. Regarding the turbulence characteristics, these simulations agree with those previously carried on for Resistive Ballooning Mode turbulence [31]. In the latter case, transport barriers were produced with an externally imposed velocity shear. In particular, a strong decrease of electric potential fluctuations is always observed, whereas the decrease of density or pressure fluctuation amplitude is small in weak barriers. Thus the level of density fluctuations is not always a good signature of ITB formation.

6. LOW ORDER RATIONAL Q MIN AND DOUBLE INTERNAL BARRIERS.

The favourable role of a low order rational value of the minimum safety factor has been long emphasised for in JET Optimised Shear plasmas [22,32]. This role has been confirmed recently in reversed shear plasmas thanks to the observation of Alfvén wave cascades [32,33]. The q profile in JET during a current ramp-up is such that q_{\min} decreases with time, crossing successively several low order rational surfaces. The case of $q_{\min} = 2$ is intriguing and analysed in detail in a companion paper [33]. An example is shown in figure 4 that shows contour lines of ρ_{T*} for the Pulse No: 51573. First a barrier appears at $R \approx 3.35\text{m}$ in a region where the shear is negative. A dramatic change of structure appears at $t \approx 6\text{s}$. This corresponds to the appearance of the surface $q=2$ at q_{\min} . Then two barriers appear that follow approximately the two $q=2$ surfaces.

A first explanation relies on MHD modes located at $q=2$ generating a localised velocity shear. A

good correlation between ITB formation and MHD activity was found in positive (optimised) shear plasmas [33]. On the other hand no strong MHD activity is observed in reversed shear plasmas apart from the Alfvén cascade itself. However tearing modes located at $q=2$ surfaces may be difficult to detect. Turbulence itself could be responsible for a flow generation close to rational q values. This explanation does receive some support from electromagnetic turbulence simulations with the CUTIE code [34]. These simulations also show that the bootstrap current is enhanced near rational q values, thus further lowering the magnetic shear locally.

A second explanation relies on the existence of gaps in the density of low wave number rational surfaces. This gap is wider when q_{\min} is close to a low order rational number. It depends sensitively on the curvature of the q profile [30]. Also gaps tend to develop in the vicinity of low order rational numbers even for finite magnetic shear. A comparison between the radial position of resonant surfaces such that $k_{\theta}\rho_{s0} < 1$ and the actual evolution of the barrier gives a remarkable agreement ([33]). First a large gap appears just before $q_{\min}=2$ (typically for $2-q_{\min}$ of the order of a few 10^{-3}). Second, once q_{\min} becomes smaller than 2, two gaps follow the $q=2$ surfaces, whereas the central gap close to q_{\min} contains high wave number resonant surfaces. It may therefore be possible that a strong barrier only appears when q_{\min} crosses the $q=2$ surface, then splits. Coexistence of barriers is possible, as shown in Fig.5. The same figure shows that the barriers are stronger near $q=2$ than near q_{\min} . Note, however, that an explanation based on the density of rational surfaces does not explain the onset and self-sustainment of a barrier located somewhat in the negative shear region (as in Pulse No: 51573 before $t=6s$). Thus both $s < 0$ and $s=0$ (and rational q_{\min}) must be invoked to explain the whole history of this kind of plasma.

7. CONCLUSION

Many experimental results in JET indicate that the onset of an ITB is sensitive to the profile of the safety factor. Part of these observations can be explained by the dependence of the linear growth rate on the safety factor and its gradient. This result is confirmed by both linear stability analysis and turbulence simulations. Models based on a transport reduction due to magnetic shear combined with velocity shear also reproduce the data in a satisfactory way. Many models fail to explain the particular role of rational surfaces. However two explanations are possible. One is based on MHD or low m,n turbulence modes generating a localised $E \leftrightarrow B$ shear flow. The second explanation relies on the development of a region without any low wave number resonant surface. Turbulence simulations confirm the possible coexistence of several barriers. They also indicate that rational q surfaces play a special role.

REFERENCES

- [1]. J.F. Drake, Y.T. Lau, P.N. Guzdar, et al., Phys. Rev. Lett. **77**, 494 (1996).
- [2]. F. Romanelli and F. Zonca, Phys. Fluids B **5**, 4081 (1993).
- [3]. R.E. Waltz, G.D. Kerbel, J. Milovitch, and G.W. Hammett, Phys. Plasmas **2**, 2408 (1995).

- [4]. H. Biglari, P. Diamond, and P.W. Terry, Phys. Fluids B **2**, 1 (1990).
- [5]. T.S. Hahm and K.H. Burrell, Phys. of Plasmas **2**, 1648 (1995).
- [6]. F.L. Hinton and G.M. Staebler, Phys. Fluids B **5**, 1281 (1993).
- [7]. G. Tresset, X. Litaudon, D. Moreau, et al., Nucl. Fusion **42**, 520 (2002)
- [8]. P. Mantica et al., this conference.
- [9]. R.C. Wolf et al., “Characterisation of ion heat conduction in JET and ASDEX-Upgrade plasmas with and without internal transport barriers”, submitted to Plasma Physics and Control Fusion.
- [10]. Y. Baranov, private communication.
- [11]. R.V. Budny, R. Andre, A. Becoulet, et al. submitted to Plasma Physics and Cont. Fusion.
- [12]. F. Crisanti, B. Esposito, C. Gormezano et al., Nucl. Fusion **41**, 883 (2000).
- [13]. L.-G. Eriksson, C. Fourment, V. Fuchs et al. , Phys. Rev. Lett. **88**, 145001 (2002).
- [14]. J. Weiland, “Collective Modes in Inhomogeneous Plasmas”, IOP, 2000.
- [15]. A.L. Rogister, Nucl. Fusion **41** 1101 (2001).
- [16]. M. Kotschenreuther, W. Dorland, M.A. Beer, et al., Phys. Plasmas **2**,2381 (1995).
- [17]. D.E. Newman, B.A. Carreras, D. Lopez-Bruna, et al., Phys. Plasmas **5**, 938 (1998).
- [18]. C. Bourdelle et al., Nucl. Fusion **42**, 892 (2002).
- [19]. W.A. Houlberg, K.C. Shaing, S.P. Hirshman, and M.C. Zarnstorff, Phys. Plasmas **4**, 3230.
- [20]. J. Kim et al., Phys. Rev. Lett. **72**, 2199 (1994).
- [21]. P. Maget , B. Esposito, E. Joffrin ,”Statistical analysis of Internal Transport Barriers in JET”, submitted to Nucl. Fusion.
- [22]. C. Challis et al., Plasma Phys. Control. Fusion **44**, 1031 (2002).
- [23]. B. Esposito et al, “JET internal transport barriers: experiment vs theory”, submitted to Plasma Physics and Control. Fusion (2002)
- [24]. T.J.J. Tala, J.A. Heikkinen, V. Parail et al., Plasma Physics and Control. Fusion **43**, 507 (2001).
- [25]. F. Imbeaux et al., EU-US TTF meeting, Cordoba 2002.
- [26]. P. Zhu, G. Bateman, A.H. Kritz and W. Horton, Phys. Plasmas **7**, 2898 (2000).
- [27]. T.J.J. Tala, V.V. Parail, A. Becoulet, et al., Plasma Physics and Control. Fusion **44**, A495 (2002).
- [28]. X. Litaudon, F. Crisanti, B. Alper, Plasma Phys. Control. Fusion **44**, 1057 (2002).
- [29]. X. Garbet, C. Bourdelle, G.T. Hoang, et al., Phys. Plasmas **8**, 2793 (2001).
- [30]. I. Voitsekhovitch et al., to appear in Phys. of Plasmas.
- [31]. P. Beyer, S. Benkadda, X. Garbet, P.H. Diamond, Phys. Rev. Lett. **85**, 4892 (2000).
- [32]. E. Joffrin et al., Plasma Physics and Control. Fusion **44**, 1739 (2002).
- [33]. E. Joffrin et al., this conference
- [34]. A. Thyagaraja, Plasma Phys. Control. Fusion **42**, B255 (2000).




Figure 1: Profiles of safety factor and ion temperature, linear growth rates and velocity shear rate of JET Pulse No: 51976 at $t=6s$.




Figure 2: Profile modelling comparison for the Pulse No: 53521.




Figure 3: Left panel: experimental profiles of safety factor, electron and ion temperature of the JET Pulse No:53521 at $t=10s$ (q profile from [28]). Right panel: turbulence simulation of a barrier with the same q profile. Circles are the positions of $k_{\theta} \rho_{s0} < 1$ resonant surfaces on the q profile.




Figure 4: Contours of ρ_{T*} of the Pulse No: 51573 (from [33]).




Figure 5: Simulation of a barrier with reversed magnetic shear and q_{min} just below 2 with the turbulence code TRB. Circles are the position of resonant surfaces on the q profile. The dashed lines are the ion and electron pressure profiles.