

EFDA-JET-CP(04)07-32

JET

R. Nazikian, B. Alper, H.L. Berk, D. Borba, C. Boswell, R. V. Budny, K. Burrell,
C.Z. Cheng, E.J. Doyle, E. Edlund, R.J. Fonck, A. Fukuyama, N.N. Gorelenkov,
C. Greenfield, D.J. Gupta, M.Ishikawa, J. Jayakumar, G.J. Kramer, Y. Kusama,
R. La Haye, G.R. McKee, W.A. Peebles, S.D. Pinches, M. Porkolab, J. Rapp,
T.L. Rhodes, S.E. Sharapov, K. Shinohara, J.A. Snipes, W. Solomon, E. Strait,
M. Takechi, M. VanZeeland, P. West, K.L. Wong, S. Wukitch, L. Zeng
and JET EFDA Contributors

Observation of Energetic Particle Driven Modes Relevant to Advanced Tokamak Regimes

Observation of Energetic Particle Driven Modes Relevant to Advanced Tokamak Regimes

R. Nazikian¹, B. Alper², H.L. Berk³, D. Borba⁴, C. Boswell⁵, R.V. Budny¹,
K. Burrell⁶, C.Z. Cheng¹, E.J. Doyle⁷, E. Edlund⁵, R.J. Fonck¹¹, A. Fukuyama⁸,
N.N. Gorelenkov¹, C. Greenfield⁶, D.J. Gupta¹¹, M.Ishikawa⁹, J. Jayakumar¹⁰,
G.J. Kramer¹, Y. Kusama⁹, R. La Haye⁶, G.R. McKee¹¹, W.A. Peebles⁷,
S.D. Pinches¹², M. Porkolab⁵, J. Rapp¹³, T.L. Rhodes⁷, S.E. Sharapov²,
K. Shinohara⁹, J.A. Snipes⁵, W. Solomon¹, E. Strait⁶, M. Takechi⁹,
M. VanZeeland⁶, P. West⁶, K.L. Wong¹, S. Wukitch⁵, L. Zeng⁷

¹Princeton Plasma Physics Lab., P.O. Box 451, Princeton NJ 08543, USA
 ²EURATOM/UKAEA Fusion Association, Culham Science Centre, Abingdon, UK
 ³Institute of Fusion Studies, University of Texas at Austin, Austin, Texas 78712-02644
 ⁴EURATOM/IST Fusion Association, Centro de Fusao Nuclear, Lisboa, Portugal
 ⁵MIT, Plasma Science & Fusion Center, 167 Albany Street, Cambridge, MA 02139 USA
 ⁶General Atomics, La Jolla, CA 92186-5608
 ⁷University of California Los Angeles, Los Angeles, California 90095-1597
 ⁸Kyoto University, Kyoto 606-8501, Japan
 ⁹Japan Atomic Energy Research Institute, Naka-machi, Ibaraki-ken, Japan
 ¹⁰Lawrence Livermore National Laboratory, Livermore, CA 94550
 ¹¹University of Wisconsin - Madison, Madison, WI 53706
 ¹²Max-Planck Institut fuer Plasmaphysik, Euratom Association, D-85748 Garching, Germany
 ¹³Institut für Plasmaphysik, Forschungszentrum Juelich GmbH, Euratom association, TEC, Jülich, Germany

(Proc.20th IAEA Fusion Energy Conference, Vilamoura, Portugal (2004).

Preprint of Paper to be submitted for publication in Proceedings of the 20th IAEA Conference, (Vilamoura, Portugal 1-6 November 2004) "This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK."

"Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK."

ABSTRACT

Measurements of high frequency oscillations in JET, JT-60U, Alcator C-MOD, DIII-D and TFTR plasmas are contributing to a new understanding of fast ion driven instabilities relevant to Advanced Tokamak (AT) regimes. A model based on the transition from a cylindrical-like frequency-chirping mode to the Toroidal Alfvn Eigenmode (TAE) has successfully encompassed many of the characteristics seen in experiments. In a surprising development, the use of internal density fluctuation diagnostics has revealed many more modes than has been detected on edge magnetic probes. A corollary discovery is the observation of modes excited by fast particles traveling well below the Alfv $\mathbf{\check{z}}$ velocity. These observations open up new opportunities for investigating a $\mathbf{\hat{\omega}}$ a of Alfv $\mathbf{\check{z}}$ Eigenmodes $\mathbf{\hat{m}}$ present scale experiments, and highlight the need for core fluctuation and fast ion measurements in a future burning plasma experiment.

1. INTRODUCTION

The physics of energetic particle driven modes is of intrinsic interest for understanding wave-particle interaction as well as of practical interest for the impact such interactions may have on the plasma. High frequency oscillations such as Alfv $\check{\mathbf{z}}$ waves can perturb resonant fast ion orbits. If the modes grow to sufficient amplitude then the fast ions may be lost to the wall or transported out of the plasma core. Understanding the stability and saturation mechanisms for these modes in present experiments is the key to predicting fast ion behavior in future burning plasma experiments.

Fusion researchers have made steady progress towards developing the Hybrid Regime and other noninductive operating regimes as a possible alternative to ELMy H-mode operation for a burning plasma experiment. In parallel, substantial progress has been made towards understanding the physics of energetic particle driven modes relevant to these advanced operating regimes. The rapid progress in understanding is due in large part to a remarkably strong collaboration between colleagues in the USA, Japan and Europe.

In this paper we present recent developments in our understanding of energetic particle driven modes in weak and reverse magnetic shear plasmas that are most relevant to Advanced Tokamak (AT) plasma regimes. Here AT refers to the development of high confinement steady state plasmas, which typically require weak and/or reverse magnetic shear. Foremost among these recent developments is the critical importance of core fluctuation diagnostics for identifying modes not readily observable on external magnetic probes. A corollary discovery is the simultaneous excitation of many fast ion driven waves with very high toroidal mode numbers. As a measure of our growing understanding, in the 2002 Snowmass report [1] there was no explicit calculation of the stability of modes in reverse magnetic shear configurations. A similar study conducted today would not be acceptable without a detailed analysis of Alfvž eigenmodes in AT relevant plasma regimes.

2. BACKGROUND

Low frequency chirping modes were initially observed in a frequency range appreciably below and up to the TAE frequency. They arose in several tokamaks such as: the TFTR-DT experiments where the modes were driven by fusion alpha particles in weak magnetic shear plasmas [2], and in several hydrogen minority Ion Cyclotron heating experiments in JT-60U, JET and Alcator C-MOD using Hydrogen minority Ion Cyclotron heating [3, 4, 5, 6]. An impressive early observation of the frequency chirping on JT-60U [3, 4] is shown in Figure 1. These modes exhibit a sequence of increasing toroidal mode number as the magnetic safety factor decreased toward an integer value and not much frequency downshifting, a characteristic of most data where this type of mode arises. In addition the higher n-modes sweep at a faster rate. A significant feature of this data is the observation of high toroidal mode numbers (up to n=11) on external magnetic probes. Another feature of all the data is that these modes occur when the equilibrium has a region of reverse magnetic shear as was clearly observed on JET [6].

Initially, the modes on TFTR were misidentified as TAEs due to the proximity of the mode frequency to the TAE frequency, with some caveats associated with low-n modes. However, the more dramatic frequency chirping observed on JT-60U, JET and Alcator C-MOD indicated a need to find an alternative Alfvžic mode. Early analytic [7] and numerical [8] studies identified modes where the real frequency was found to be sensitive to the evolution of qmin. A detailed theory-experiment comparison on JET [7, 9] and JT-60U [10] convincingly demonstrates that these chirping modes are indeed Alfvž Eigenmodes with a frequency determined predominantly by the evolution of q_{min} . A subsequent reevaluation of the TFTR data resolved earlier inconsistencies with the TAE theory [11]. These developments mark dramatic progress in providing a uniform understanding of these curious frequency chirping modes in reverse magnetic shear plasmas in a wide range of conditions and machines.

The basic picture that has emerged from these studies is that cylindrical-like Alfvž modes (i.e., modes with one dominant poloidal mode number m) appear at rational values of $q_{min} = m/n$. The modes then transform to TAEs as $q_{min} \rightarrow qTAE = (m-1/2)/n$. Over this relatively small change in qvalue, the mode frequency changes from significantly below the TAE frequency to the TAE frequency. This mode is referred to as a Reverse Shear Alfvn Eigenmode (RSAE) on JT-60U and as a Cascade mode on JET, where both terms refer to the existence of modes with a single dominant poloidal harmonic m at the location of q_{min} . In the range m/n < q_{min} < (m-1/2)/n, the mode frequency is given by $\omega \approx k_{\parallel} V_A$, where $k_{\parallel} = (m - nq_{\min})/q_{\min}R$ for one dominant poloidal harmonic m and V_A is the Alfv $\mathbf{\check{Z}}$ velocity. For q _{min} > (m-1/2)/n the mode approaches the TAE frequency where it fades away (see Fig.1) or merges with the TAE as best seen in Figure 4. The rate of frequency chirping vs q_{min} is proportional to the toroidal mode number n ($\partial \omega / \partial q_{\min} \approx n V_A / q_{\min} R$) so that the mode numbers may be extracted without the use of toroidal correlation analysis. This is particularly useful for the interpretation of core interferometer measurements when external magnetic data is not available. Outstanding issues still exist such as the precise role of energetic particles [9] and the range of applicability of the ideal MHD description [12]. However these issues do not obscure the fundamental similarity of the model predictions in terms of the observable characteristics of the modes. Further progress and new insights will undoubtedly arise from more detailed measurements and calculations.

3. INTERNAL MODE MEASUREMENTS

A unique contribution of the TFTR-DT experiments is the measurement of the internal structure of the Cascade modes with the use of reflectometry [11]. The observation of an antiballooning density mode structure for the low frequency n=2 mode has been highly controversial for some time. The issue was resolved when the NOVA-K code confirmed that the Alfvž Eigenmode near q_{min} should exhibit density peaking on the high field side of the magnetic axis (Figure 2). The magnitude of the magnetic component of the calculated eigenmode is close to in-out symmetric, consistent with a single dominant poloidal mode number m.

In a recent experiment on JET, new observations of a broad spectrum of Cascade modes were observed in reverse magnetic shear plasmas by operating the plasma density below the O-mode reflectometer cutoff density (Fig.3) [13]. The interferometric data revealed a rich Alfvž spectrum consisting of many frequency-sweeping discrete modes ranging from 40 kHz to the TAE frequency \approx 140 kHz. These observations are in agreement with the established characteristics of Cascade modes driven by high-energy hydrogen minority ions heated by the Fast Waves. The new evidence from the plasma core indicates that Cascades are far more prevalent in AT relevant regimes and in higher numbers than previously thought, with mode numbers up to n = 16 observed in the plasma core. These mode numbers are not directly measured but inferred from the rate of change of frequency of the modes. Note that these observations are in contrast to isolated low-n modes observed on external magnetic probes. As a q-profile diagnostic, the onset of the Grand Cascade (dashed line in Figure 3) accurately identifies the entry of $q_{min} = 2$ surface in the plasma. The detection of core localized high-n modes allows a more accurate determination of q_{min} than is possible with external magnetic measurements. It also allows for a more accurate study of the stability of these modes under varying plasma conditions and auxiliary heating.

In a parallel development, the Phase Contrast Imaging (PCI) diagnostic on Alcator C-MOD has been highly successful in identifying new mode activity not observable on external magnetic probes [14] (Fig.4). The excitation of Alfvž eigenmodes has been observed in Alcator C-MOD with hydrogen tail ion temperatures as low as 100keV from TRANSP analysis. Both in JET and in Alcator C-MOD, core chirping modes are observed that cannot be identified on external magnetic probes. Due to the dominant Radio Frequency (RF) heating in these experiments, the effect of toroidal rotation on the mode frequency is weak compared to the dependence on q_{min}. The modes observed on interferometer channels typically chirp faster than modes observed on magnetic diagnostics, indicating that higher toroidal mode numbers are involved. The work on Alcator C-MOD and also recent data obtained on DIII-D demonstrates that CO₂ laser interferometry is sufficiently sensitive to detect these modes in fusion plasmas. On JET the interferometer measurements in Figure 3 are taken with the O-mode reflectometer system where the density of the plasma is kept just below the O-mode cutoff. In a similar approach, line integral density fluctuation measurements were taken with an X-mode reflectometer system on JT-60U. For a 1 T magnetic field, the launch frequency (≈140GHz) propagates about the third electron cyclotron harmonic and reflects from the inner wall. Figure 5 shows a comparison of magnetic and line density measurements

in a 1 T discharge where the modes are excited by the injection of 360 keV neutral beam ions. Note that the mode amplitude peaks earlier in the line density than on the edge magnetic signal for Mode-1. This is similar to the behavior of core reflectometer measurements on TFTR [11]. These observations are particularly important when attempting to correlate fast ion loss with mode amplitude measurements.

The many core-localized modes not observable on magnetic sensors in JET highlight the need for core density fluctuation diagnostics on a future burning plasma experiment.

4. NEUTRAL BEAM DRIVEN ALFVÉN EIGENMODES

Extensive studies have been conducted on JT-60U using tangentially injected neutral beams with energies in the range 360 keV and toroidal magnetic field near 1 T where the beam ion velocity is close to the Alfv $\mathbf{\check{z}}$ speed. However, recent studies have shown that RSAEs or Cascades can be excited at much higher toroidal magnetic fields on JT-60U where the beam ion velocity is closer to half the Alfv $\mathbf{\check{z}}$ speed.

These observations highlight a general misconception that high-energy ions are required to excite Alfvž Eigenmodes in present scale experiments, and that these energies exceed the range available in standard positive ion heating beams unless the toroidal magnetic field is greatly reduced. Recent observations on JET and DIIID challenge this notion and so we need to consider carefully the resonance condition for these modes.

For passing particles the resonance condition for the mode-particle interaction is given by $k_{\parallel}VA$ = $(k_{\parallel} \pm p/qR)V_{\parallel}$ where p is a positive integer and $k_{\parallel} = (m - nq)/qR$. Hence resonance is at $V_{\parallel}/V_A = 1/2$ $[1\pm 2p/a]$ where $\alpha = \omega_m/\omega_{TAE}$, $\alpha_{TAE} = V_A/2qR$ and $\omega_m = k_{\parallel}VA$. As q varies from m/n < q < (m D/m)2) /n, a single Cascade mode with frequency ω_m sweeps up to the TAE frequency ω_{TAE} . Thin orbits have the strongest interaction for the p=1 and p=2 cases which gives for $\alpha \approx 1$ (mode near the TAE frequency) the resonance condition $|V_{\parallel}/V_{A}| \approx 1$ or 1/3. Note that for even for p = 1 and for a Cascade mode with small α corresponding to a frequency well below the TAE frequency, we obtain $|V_{\parallel}/V_{A}| \approx \alpha / (2 \text{ D}\alpha)$. In experimental observations the minimum of the Cascade frequency has been observed as low as ~0.3 ω_{TAE} , so that the Cascade p = 1 resonance can occur below the TAE sideband resonance condition with sub- Alfvžic beams. Then the mode would fade away as the frequency increases because the resonance condition cannot be fulfilled at higher frequency. This is possibly occurring in the data of Figure 7 on JET. However, if the p=2 resonance condition is excited (where $|V_{\parallel}/V_{A}| = 1/3$) in sub-Alfvžic beam experiments, then the resonance condition is more likely to be satisfied over the entire Cascade frequency band. Even lower velocity beams can achieve resonances at higher p values, but these higher order resonances require finite size orbit widths (orbit width larger than $\epsilon r/n$) to obtain a substantial interaction. Further, beams are often partially injected into the trapping region and Coulomb scattering causes an accumulation of trapped particles, where the precession drift, ω_D , is significant. Then, as $k_{\parallel}V_A = (k_{\parallel} \pm p/qR)V_{\parallel} + n\omega_D$, is the resonance condition, the precessional drifts may enable resonance to be achieved at yet lower beam velocities.

Recent FIR scattering measurements on DIII-D have revealed clear evidence for the excitation of Cascade modes in weakly reverse magnetic shear plasmas in 2.1 T plasmas with 80 keV beams. The Far Infra Red (FIR) scattering system detected fluctuations in the range $D < k_{\theta} < 1 \text{ cm}^{-1}$. These modes have since been observed in many DIII-D plasmas with weak and reverse magnetic shear. A notable characteristic of the observations on DIII-D is the lack of correlation between core density and edge magnetic oscillations. Figure 6 shows an example of a Grand Cascade observed on DIII-D with the FIR scattering system. These are concentrations of Cascade modes that indicate integer q_{min} crossings. Note that the observed modes exhibit very strong bursting behavior on the scattering signal, suggesting a strong interaction and possible redistribution of the beam ions. Comparison with MSE measurements in different plasma discharges confirms that the Grand Cascades on DIII-D correspond to integer qmin crossings. The mode activity commences early in the beamheatingphase when the deuterium beam ion velocity at 80keV is less than 40% of the AlfvŽ velocity. It is difficult on DIII-D to inject beam ions well below the sideband resonance condition for TAEs ($V_{\parallel} = V_A/3$). However, it is quite possible that positive ion beam injection below the sideband resonance condition can be achieved on JT-60U and JET.

Evidence from JET indicates that Cascade modes can also be excited by neutral beam injection in high toroidal field plasmas. In this plasma LHCD was used to generate the current hole but no ICRH was used. Figure 7 shows data from an O-mode interferometer channel with 2.3MW deuterium beam injection (110keV) in a 3.46T plasma with central electron density 2.0×10^{13} cm⁻³. The modes were observed in a current hole discharge where the q-profile is strongly reversed and the time of the Grand Cascade corresponds to the crossing of $q_{min} = 3$. The neutral beam ion velocity is ≈ 0.27 V_A . However the angle of neutral beam injection in this case is 45 degrees to the field lines so that collisions during the slowing down process can produce a significant population of trapped energetic ions. A detailed analysis of the resonance condition is needed to determine the excitation mechanism for these modes and the relative role of trapped vs passing particles. However, what is clear is that interferometer measurements have succeeded to identify many Cascade modes driven by energetic ions traveling close to or below $V_A/3$.

Note that with neutral beam injection there is now significant plasma rotation. Consequently, there is a Doppler shift of the spectrum to higher frequency with increasing toroidal mode number as shown in Figure 7. Higher mode numbers are expected when approaching integer q_{min} values from above. This should form an envelope of modes with increasing frequency due to the Doppler shift. Once q_{min} drops below 3 the sequence of modes reverses toward low-n. Further work is still needed to fully explore the resonance condition for Cascade modes in JET either by raising the toroidal field strength or by lowering the beam ion voltage.

Evidence for the excitation of many high-n modes is observed on the DIII-D facility. For a long time it has been speculated that high frequency modes could be responsible for the anomalous beam ion diffusion inferred (not measured) in weak and reverse magnetic shear plasmas on DIII-D [15]. The observed modes typically extended up to 1MHz and were most clearly seen on FIR scattering signals, with no clear counterpart on magnetic diagnostics.

Figure 8 shows an example of the mode activity observed in a Quiescent-H-mode (QH) plasma with weakly reversed magnetic shear, together with a simple model analysis based on the observed evolution of q_{min} from MSE measurements. The modeling assumes a transition from the Cascade mode (starting at $q_{min} = m/n$) to the TAE [at $q_{min} = (m-1/2)/n$] for a wide range of toroidal mode numbers, and adds a Doppler shift using measured charge exchange data. In this discharge the neutral beams are injected in the direction opposite to the plasma current and the observed mode activity is consistent with mode propagation in the direction of beam injection in the plasma rest frame. More recent data analysis has indicates that the modes in the plasma rest frame are propagating opposite to the direction of the plasma current and in the direction of neutral beam injection. (This observation introduces a puzzle as the theoretically expected direction of mode propagation for this case is in the direction of the plasma current in the frame where the plasma is at rest with respect to the mode location). The different bands of modes correspond to different poloidal mode numbers m = n + 1 where l = 0, 1, 2 ... correspond to increasing frequency bands. For each of the three dominant bands of modes we obtain an average frequency spacing of 33 kHz, 36 kHz and 33 kHz (for the l = 0, 1, and 2 bands respectively). The central toroidal rotation in this discharge and at this time is 33 kHz, while at the location of q_{min} it is 30 kHz. The magnitude of the frequency shift is certainly consistent with that expected for successive toroidal mode numbers n, n + 1, fMode numbers up to n = 30 are inferred from the Doppler shifted spectra with up to 20 independent modes observed simultaneously in some discharges. If we now look back to Figure 3, we can see just such a similar set of nested bands on JET. However unlike in DIII-D, the bands of modes on JET are partially obscured by low plasma rotation. In the DIII-D data the plasma is rotating at $\approx 5\%$ of the Alfvž velocity in the direction of neutral beam injection.

In Figure 8 the highest mode numbers can have very short poloidal scale lengths. Note that for m = n = 30 and r \approx 20 cm in DIII-D the poloidal scale length can be in the range k₀ \approx 1 cm⁻¹ so that a small angle scattering system turns out to be quite effective for the investigation of these high-n modes. Also, with such core localized high mode numbers it is unlikely that there will be a detectible magnetic component to these modes at the plasma edge. Recent experiments have corroborated the FIR measurements on DIIID with extensive reflectometer, vertical and radial CO₂ interferometer and Beam Emission Spectroscopy measurements. The modeling reproduces the qualitative behavior of the data, however a quantitative description must await a detailed stability analysis using NOVA-K.

5. RELEVANCE TO BURNING PLASMAS

The excitation of Alfvž eigenmodes with low energy heating ions raises the new possibility of exploring a $\hat{\omega}a$ of modes $\hat{\omega}$ which has been predicted to occur in a burning plasma experiment. For similar MHD equilibria, the maximum toroidal mode number excited in a device should scale as the ratio of the average minor radius to the fast ion gyroradius, $n_{max} \approx a/(q \rho_{\alpha})$ where ρ_{α} is the Larmor radius of the alpha particles. In addition the upper limit for the number of excited modes should scale as ϵn_{max}^2 . The higher the mode number n_{max} the more modes can potentially be excited. In ITER the ratio is of the order 45-50 for 3.5 MeV super Alfvžic alpha particles. For 80

keV deuterium ions in 2.1 T DIII-D plasmas the ratio is ≈ 30 for $V_{\parallel}/V_A \approx 0.3$, while for 110 keV deuterium ions in a 3.5 T plasmas on JET the ratio is close to 70 near the sideband resonance condition $V_A/3$.

The investigation of a Sea of Alfvž Eigenmodes In present day devices should be pursued for what it can teach us about multimode interactions. Understanding instabilities driven by low energy heating ions in existing facilities is highly relevant to the development of the Advanced Tokamak (AT) concept as well as to burning plasma science where the condition of multimode interactions is considered particularly important. The availability of highly sensitive core fluctuation measurements represents a major breakthrough for the quantitative analysis of fast ion collective instabilities. In order to fully explore this exciting area of fusion science, facilities will need a comprehensive set of core fluctuation measurements together with effective means of measuring fast ion redistribution and loss. Such a comprehensive diagnostic set may turn out to be a necessity in a future burning plasma experiment.

ACKNOWLEDGMENTS

This work is supported by DOE contracts DE-FC02-04ER54698, DE-AC02-76CH03073 and DOE Grant No. DE-FG03-01ER54615. The work is also partially funded by EURATOM, the UK Engineering and Physical Sciences Research Council, and is partly performed under the European Fusion Development Agreement (EFDA). The work is also partially funded by the Japan Atomic Energy Research Institute and is implemented under the International Energy Agency is co-operation agreement on Large Tokamak Facilities established between JAERI, DOE and EURATOM, and by the US-EU bilateral agreement on fusion energy established between DOE and EURATOM.

REFERENCES

- [1]. Snowmass Fusion Summer Study, 2002
- [2]. R. Nazikian et al., Proceedings of the 16th IAEA International Conference on Fusion Energy, Montreal, 7-11 October 1996, IAEA-CN-64/A2-4
- [3]. H. Kimura, et al., Nuclear Fusion **38** 1303 (1998)
- [4]. Y. Kusama et al., Nuclear Fusion **38** 1215 (1998)
- [5]. J. A. Snipes et al., Plasma Physics and Controlled Fusion 42 381 (2000)
- [6]. S. E. Sharapov et al., Phys. Lett. A289 127 (2001)
- [7]. H.L. Berk et al., Phys. Rev. Lett., 87 185002 (2001)
- [8]. A. Fukuyama et al., Proceedings of the 5th IAEA TCM on Alpha Particles in Fusion Research, 8-11 Sept. 1997, Abingdon, UK (IAEA, Vienna 1997) [see also A. Fukuyama et al, Binetic Global Analysis of Alfvž Eigenmodes in Toroidal PlasmasOn the Proceedings of the 19th IAEA Fusion Energy Conference, Lyon, France 2002. TH/P3-14]
- [9]. S.E. Sharapov et al., Physics of Plasmas 9 2027 (2002)
- [10]. M. Takechi et al., Proceedings of the 19th IAEA Fusion Energy Conference, Lyon, France 2002 [see also Shinohara et al., NF 41 603 (2001)]
- [11]. R. Nazikian et al., Phys. Rev. Lett. 91 125003 (2003)

- [12]. G. J. Kramer et al., Plasma Phys. Control. Fusion 46 L23-L29 (2004)
- [13]. S. E. Sharapov et al., Phys. Rev. Lett. 93 165001 (2004)
- [14]. J. Snipes et al., EPS 2004
- [15]. C. M. Greenfield et al., Plasma Phys. Control. Fusion 44 A123-A135 (2002)



Figure 1: Temporal evolution of Alfvén mode frequency changes observed in a JT-60U ICRF heated RS plasma [3].

Figure 2: (a) Measured radial structure of n=2 density fluctuations along midplane on TFTR reveals agreement with Cascade eigenmode from NOVA-K. (b) Corresponding magnetic field from NOVA-K is consistent with a cylindrical-like mode.



Figure 3: Amplitude of plasma perturbations as function of time and frequency measured in JET discharge. (top) Interferometer measurements with microwave beam of 45.21GHz along plasma midplane showing up to n=16 modes and (bottom) measurements with external magnetic pick-up coils with n=3-5.



Figure 4: Core Alfvén Cascades on Phase Contrast Imaging (PCI) on Alcator C-MOD have higher mode numbers than the more global AC's seen on edge magnetic pickup coils. The PCI chords traverse the central region of the plasma where magnetic shear reversal is predicted.



Figure 5: (a) Magnetic fluctuations peak at the transition form the slow frequency-chirping mode (Mode-1) to the TAE in JT-60U. (b) Interferometer measurements identify additional low frequency chirping mode (Mode-2). The peak of the interferometer signal occurs at low frequencies, contrary to the magnetic signals, as observed on TFTR.



Figure 6: RSAEs excited with 80keV tangential beam injection in a DIII-D discharge with $B_T = 1.8$ T, R = 1.7m and $n_e(0) 2.0 \times 10^{19}$ cm⁻³. The Spectrogram of density fluctuations is from FIR scattering measurements through the plasma midplane. The magnetic probes have an effective bandwidth of 250kHz. Note that strong bursting and fast chirping behavior of the mode amplitude observed at low frequency indicates a strong interaction with neutral beam ions.



Figure 7: Spectrogram of Cascade modes excited by 110keV tangentially injected deuterium beam ions in JET: $B_T = 3.46 \text{ T}, n(0) = 2.0 \times 10^{13} \text{ cm}^{-3}, R = 300 \text{ cm}.$



Figure 8: (a) The time evolution of a number of frequency bands in the density fluctuation spectrum from FIR scattering measurements in DIII-D ($0 < k_{\theta} < 2cm^{-1}$). Modes are excited by 80keV deuterium beam ions. Plasma parameters: $B_T = 2.1T$, $n_e(0) = 2.6 \times 10^{13} \text{ cm}^{-3}$, R = 1.7m, tor33kHz. (b) Model analysis of Cascade modes transitioning to TAEs based on evolution of qmin from MSE measurements. In this data range it is assumed that qmin varies linearly with time.