
EFDA–JET–CP(07)03/24

T.M. Biewer, Y. Andrew, R.E. Bell, K. Crombe, C. Giroud, N.C. Hawkes,
D.L. Hillis, A.G. Meigs, C.R. Negus, A.D. Whiteford, K.-D. Zastrow
and JET EFDA contributors

Charge Exchange Recombination Spectroscopy Measurements from Multiple Ion Species on JET

"This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK."

"Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK."

Charge Exchange Recombination Spectroscopy Measurements from Multiple Ion Species on JET

T.M. Biewer¹, Y. Andrew², R.E. Bell³, K. Crombe⁴, C. Giroud², N.C. Hawkes²,
D.L. Hillis¹, A.G. Meigs², C.R. Negus², A.D. Whiteford⁵, K.-D. Zastrow²
and JET EFDA contributors*

¹*Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA*

²*EURATOM/UKAEA Fusion Assoc., Culham Science Center, Abingdon, OX14 3DB, UK*

³*Princeton Plasma Physics Laboratory, Princeton, NJ 08543, USA*

⁴*EURATOM-Belgian State, Department of Applied Physics, Ghent University, Rozier 44 B-9000, Ghent, Belgium*

⁵*Department of Physics, University of Strathclyde, Glasgow, G4 0NG, UK*

** See annex of M.L. Watkins et al, "Overview of JET Results",
(Proc. 21st IAEA Fusion Energy Conference, Chengdu, China (2006)).*

Preprint of Paper to be submitted for publication in Proceedings of the
34th EPS Conference on Plasma Physics,
(Warsaw, Poland 2nd - 6th July 2007)

A programmatic goal of the Joint European Torus (JET) facility in 2009 is to implement an ITER-like wall, which implies a significant reduction in the amount of carbon present in JET discharges. As a result, carbon Charge Exchange Recombination Spectroscopy (CXRS) measurements (currently the preferred ion temperature and rotation measurement on JET) may be hindered. The current CXRS system on JET consists of a suite of instruments [1, 2, 3] simultaneously observing multiple ion species. CXRS measurements of ion temperature and rotation (toroidal and poloidal) are made (in the core^{1,2} and at the plasma edge³) both on typical ion species, such as carbon, beryllium, and helium (C, Be, He), as well as on puffed impurity ion species, such as argon, neon, and nitrogen (Ar, Ne, N). A detailed comparison between simultaneous CXRS measurements of various ion species is presented. Results indicate that the toroidal ion temperatures (T_i) and rotations (v_T) measured from C, Ne, Ar, and N are in agreement to within instrumental errors. He/Be measurements have proven difficult because of the complexity of the spectra in the 468.5 nm region, though the results are not inconsistent with the measurements from heavier impurity ions. The effect of CX ion “plumes [4, 5, 6]” is being investigated as a confounding factor in the He/Be spectra. The implementation of an ITER-like wall, and subsequent reduction in carbon edge impurity lines, may simplify the complexity of the He/Be spectra, especially if the concentration of Be in JET plasmas increases. Results include analysis using the new CXSFIT routine [7], developed to standardize spectral line fitting on JET and other plasma devices, such as ASDEX-Upgrade.

OBSERVATIONS

When attempting to assess how well two measured values agree, there are a number of statistical markers that can be used. The “similarity” of two curves (**A** and **B**) can be defined [8] as the normalized dot product: $S \equiv |\mathbf{A} \cdot \mathbf{B}| / (|\mathbf{A}| |\mathbf{B}|)$. For radial profiles (e.g. T_i , v_T) from different instruments the profile curves are made into “vectors” by interpolation onto a common radial grid, where the radius values are used as a set of basis vectors. Offsets in the measured quantities due to possible calibration errors are removed by the normalization, allowing a comparison of the profile shape (between two instruments) to proceed. By definition, $0 \leq S \leq 1$, and for typical radial profiles of CXRS measurements, $S \approx 1$. For this reason, it is useful to define the “dissimilarity:” $D \equiv 100\% (1 - S)$, which can be thought of as the percent difference between two radial profile curves.

The CXRS instruments on JET typically produce data at 100Hz, i.e. with an integration period on the order of 10ms. Each JET pulse yields between 500 and 1000 frames of data for a given CXRS instrument. Figure 1 shows a comparison between two CXRS instruments (KS5C: tuneable; KS5D: fixed wavelength), both set to measure the 529.0nm C VI CX line. “By eye” it is clear from Fig.1(a) that T_i and v_T measurements from these two instruments are in good agreement. Not surprisingly, the “dissimilarity” for these instruments (Fig. 1b, c) are $D_{T_i} = 0.064 \pm 0.002\%$, and $D_{v_T} = 0.524 \pm 0.003\%$.

Since the KS5C instrument is tuneable, it is possible to compare the CX measurements of other ion species to the C VI CX measurements of the KS5D instrument. This is shown in Figure 2 for C,

N, Ar, C/Ar, C/Ne, and He/Be. In practice some CX lines within the same spectra are fit with “coupled” T_i , νT . Hence the “dissimilarity” is a type of hybrid comparison in those cases. This is done when the CX radiance of a particular ion species is very low, e.g. for a “lightly puffed” impurity ion such as Ar or Ne.

A similar type of comparison can be done between other instruments, such as the edge CXRS, which has channels overlapping with core CXRS views. Using the propagated measurement errors and photon flux as weighting, the weighted-average D (and S) are shown in Table 1 for the various ion species, as compared to C VI.

DISCUSSION

Overall, there is a high level of agreement between the multiple measurements of T_i and νT in JET plasmas. Across instruments and across ion species, $S \sim 0.99$, compared to the theoretical maximum of $S=1$. Nevertheless, this analysis suggests that there could be resolvable (though small) differences between ion species at high CX signal levels.

ACKNOWLEDGEMENTS

This work, carried out under the European Fusion Development Agreement, supported by the European Communities and the U.S. Dept. of Energy, has been carried out within the Contract of Association between EURATOM and the U.S.D.O.E, contract DEAC05-00OR22725.


REFERENCES

- [1]. C.R. Negus, C. Giroud, A.G. Meigs, K.-D. Zastrow, D.L. Hillis, Rev. Sci. Inst. **77** (1), 10F102 (2006.)
- [2]. D.L. Hillis, D.T. Fehling, R.E. Bell, D.W. Johnson, *et al.*, Rev. Sci. Inst. **75** (10), 3449 (2004).
- [3]. Y. Andrew, N.C. Hawkes, K. Crombe, Rev. Sci. Inst. **77**, 10E913 (2006).
- [4]. R.E. Bell, Rev. Sci. Inst. **77**, 10E902 (2006).
- [5]. U. Gerstel, L. Horton, H.P. Summers, M. von Hellerman, B. Wolle, Plasma Phys. Controlled Fusion **39**, 737 (1997).
- [6]. M. Tunklev, PhD thesis, Lund University, 1999.
- [7]. A.D. Whiteford, *et al.*, “CXSFIT User Manual,” <http://adas.phys.strath.ac.uk/cxsfit>
- [8]. J. Vega, “Data retrieval based on physical criteria,” presented at Culham Science Center, Abingdon, UK 2007.

KS5D C:	D_{Ti} (%)	S_{Ti}	D_{vT} (%)	S_{vT}
KS5C C	0.12 ± 0.04	0.9988	1.89 ± 0.07	0.9811
KS5C C/Ar	0.20 ± 0.05	0.9980	0.72 ± 0.09	0.9928
KS5C C/Ne	0.76 ± 0.05	0.9924	4.03 ± 0.12	0.9597
KS5C Ar	15	0.85	34	0.66
KS5C N	2.28 ± 0.04	0.9772	1.14 ± 0.02	0.9886
KS5C He/Be	1.23 ± 0.02	0.9877	3.62 ± 0.10	0.9735
KS7A C	0.69 ± 0.14	0.9931	5.84 ± 0.98	0.9416
KS7C C	0.131 ± 0.003	0.99869	2.74 ± 0.01	0.9726

JG07.2018-3c

Table 1: CX line brightness weighted average of D and S between C and other ions for Ti and vT measurements analyzed with CXSFIT in JET discharges. The KS5 instruments cover $\sim 0 < r/a < \sim 0.9$, while KS7A covers $\sim 0.35 < r/a < \sim 0.75$ and KS7C covers $\sim 0.75 < r/a < \sim 1$.



JG07.2018-1c

Figure 1: Comparison between three JET CXRS instruments, showing (a) profile measurements of T_i , v_T and photon flux from C_{vT} . (b) The time evolution of the ‘‘dissimilarity’’ of the two core instruments, which observe the same NBI PINIs, but from opposite toroidal directions. (c) A more detailed time history, showing that D_{vT} has a correlation with the occurrence of large Type-I ELMs.




Figure 2: Comparison between two JET CXRS instruments, one fixed on the C VI 529.0nm line, showing D_{Tl} and D_{VT} variation among ion species. The horizontal axis is the product of the measured CX line radiances. (a) The scatter in D decreases dramatically when the photon flux increases. (b) Same data as in Fig.2a, but on an expanded scale to show the variation in D among ion species. Error bars have been suppressed for the sake of clarity.