EFDA–JET–CP(09)06/28

J. Bernardo, Y. Andrew, K. Crombé, S. Reyes Cortes, G. Saibene, T.M. Biewer,
J. Ferreira, N.C. Hawkes, I. Jenkins, E. de la Luna, D. McDonald, I. Nunes,
A. Salmi and JET EFDA contributors

Highly Spatially Resolved Measurements of JET Edge Toroidal Rotation in Type-I ELMy H-mode Plasmas

"This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK."

"Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK."

Highly Spatially Resolved Measurements of JET Edge Toroidal Rotation in Type-I ELMy H-mode Plasmas

J. Bernardo[1], Y. Andrew[2], K. Crombé[3], S. Reyes Cortes[1], G. Saibene[4], T.M. Biewer[5], J. Ferreira[1], N.C. Hawkes[2], I. Jenkins[2], E. de la Luna[6], D. McDonald[2], I. Nunes[1], A. Salmiand[7] and JET EFDA contributors*

JET-EFDA, Culham Science Centre, OX14 3DB, Abingdon, UK

[1]Associação EURATOM-IST, Instituto de Plasmas e Fusão Nuclear, 1049-001 Lisbon, Portugal
[2]EURATOM-UKAEA Fusion Association, Culham Science Centre, OX14 3DB, Abingdon, OXON, UK
[3]Association EURATOM-Belgian State, Department of Applied Physics Ghent University, B-9000 Ghent, Belgium
[4]FUSION FOR ENERGY, Joint Undertaking, 08019 Barcelona, Spain
[5]Oak Ridge National Laboratory, Oak Ridge, TN 37831 USA
[6]Asociación EURATOM-CIEMAT, ES Avenida Complutense 22, E-28040 Madrid, Spain
[7]Association EURATOM-Tekes, Helsinki University of Technology, PO Box 4100, 02015 TKK, Finland
** See annex of F. Romanelli et al, "Overview of JET Results",*
(Proc. 22 [nd] IAEA Fusion Energy Conference, Geneva, Switzerland (2008)).

Preprint of Paper to be submitted for publication in Proceedings of the
36th EPS Conference on Plasma Physics, Sofia, Bulgaria.
(29th June 2009 - 3rd July 2009)

1. INTRODUCTION

Type-I ELMy H-mode operation have been pointed out as a reference operational scenario for ITER (Q = 10) [1]. However, the heat and particle flux produced by Type-I ELMs would lead to a rapid erosion of the divertor plates. In this way, several approaches have been made to control Type-I ELMs, such as in DIII-D [2], in TCV [3] and in ASDEX-Upgrade [4]. Reports from JT-60U [5] have shown that Neutral Beam Injection (NBI) in counter-current direction alters the toroidal velocity (v_φ) decreasing the frequency of Edge Localized Modes (ELMs). Turbulence is thought to be suppressed by **E**×**B** shear for which rotation is a key parameter. As a result, plasma rotation may have a direct effect on the overall plasma performance. Therefore further understanding of ELM behavior and the relationship with ripple and toroidal velocity is relevant for ITER.

Due to the finite number of Toroidal Field (TF) coils the magnetic field varies toroidally. The TF ripple amplitude (δ) is defined as the variation of the magnetic field, $\delta = (B_{max} - B_{min}) / (B_{max} + B_{min})$. The values shown in this paper are the maximum values at the plasma separatrix. For standard JET operation at midplane $\delta = 0.08\%$, while for ITER it will be of $\delta = 0.5\%$ with the inclusion in the design of Ferritic Insets (FI) compensation. JET tokamak has 32 TF coils, which can be fed with equal current or powered independently with different current in the odd and even-numbered coils, allowing to vary TF ripple amplitude up to $\delta \sim 3\%$. The TF ripple breaks the axisymmetry of the magnetic field, enhancing particles losses of fast and thermal ions [6,7] and affects the toroidal rotation velocity (v_φ).

In this paper the effect of the TF ripple on localized measurements of edge plasma toroidal rotation, close to the top of the ion temperature pedestal (T_i^{ped}), and its influence on ELM frequency is examined on JET Type-I ELMy H-mode plasmas.

2. EXPERIMENTAL RESULTS

The pulses used in this paper were performed at JET, in 2007 and 2009, in dedicated experimental campaigns for ripple studies. The following analysis has been performed for several Type-I ELMy H-mode plasmas with varying amounts of TF ripple (from the standard JET ripple $\delta = 0.08\%$ up to a maximum of 1%) and edge plasma densities. The set of pulses was carried with a plasma current from 2.6MA down to 2.4MA, while the strength of the toroidal magnetic field ranged from 2.2 T to 2.4T. In all cases, additional heating was provided by co-current NBI input power ranging from 8.6 to 15.9MW, and for some pulses Ion Cyclotron Resonant frequency Heating (IRCH) was applied between 0.7 and 1.5MW.

Detailed measurements of the edge plasma toroidal rotation velocity ($v\Delta$) and ion temperature (Ti) have been carried out using the edge Charge eXchange Recombination Spectroscopy (CXRS) diagnostic system at JET, with a temporal resolution of 50ms and a spatial resolution up to 2cm [8]. The measured quantities are those of the fully ionized carbon impurities at wavelength $\lambda = 529.1$ nm. It is assumed in the paper that the deuterium ions have the same velocity and temperature.

Figure 1 shows time traces of plasma current a-b), magnetic field c-d), volume averaged electron

density e-f), diamagnetic energy g-h), D_α emission on the outer divertor i-j) and additional heating power k-l) for Pulse No's: 77108 ($\delta = 0.08\%$) and 77110 ($\delta = 0.75\%$).

For each of the analyzed pulse several time windows of 0.5s were selected, represented by the red boxes in figure 1. During each time window the ELM frequency (f_{ELM}) was determined, using the D_α signal of the outer divertor. Toroidal velocity at the top of the pedestal (v_φ^{ped}) has been averaged over the selected time windows. The ion temperature profiles (figure 2a), given by the edge CXRS, were used to determine the T_i^{ped} location, allowing to retrieve, at the same radial position, v_φ^{ped} from the toroidal velocity profiles given by the edge CXRS diagnostic (figure 2b).

Figure 3b) show the results for pulses with higher edge density and collisionality ($\nu^* \sim 0.5$). In this set of pulses $\bar{v}_\varphi^{ped}$ ranges from -20 to +8km/s. A similar relationship between the f_{ELM} and $\bar{v}_\varphi^{ped}$ has been found, fELM increases with decreasing v , however the values of toroidal rotation velocity are considerably lower than for the low density cases.

The effect of the density on the edge rotation has been investigated further. As an example two Pulse No's: 69837 ($n_e^{ped} \sim 3.6\times10^{19}\,m^{-3}$) and 69858 ($n_e^{ped} \sim 6.1\times10^{19}\,m^{-3}$) with $\bar{v}_\varphi^{ped}$ of -100 and -9 km/s respectively are compared. Both pulses have ´=0.08% and are represented in black dots in figure 3.

Figure 4a) shows the angular momentum density profile (LΔ) for both pulses.

It can be seen that L_φ is lower over the whole profile for the high density pulse, which implies that v_φ for Pulse No's: 69858 must be considerably lower than for Pulse No: 69837, which is consistent with the obtain values for $\bar{v}_\varphi^{ped}$, which are of -100 and -9 km/s respectively.

In both pulses, the same PINIs configuration was used. The difference in total torque applied to the plasma is less than 1%. The torque profiles calculated by the neutral beam deposition code PENCIL are compared in figure 4b). For the Pulse No: 69837 the torque profile given by the orbit following Monte Carlo code ASCOT [9] is also shown and agrees with Pencil calculation. A higher edge density leads to a higher beam deposition at the edge and lower deposition in the core, as it can be seen in figure 4b). The slightly higher torque at the edge of the plasma for the higher density case (Pulse No: 69858) might enhance slightly the edge rotation, but the large difference in mass results in lower v values.

CONCLUSIONS AND DISCUSSION

The JET ripple experiments clearly show that increasing TF ripple lead to counter rotation at the plasma edge due to fast and thermal ion losses. It has been found that the toroidal velocity at the pedestal location and the ELM frequency are related. As the toroidal velocity approaches values close to zero, for TF ripple of $\leftrightarrow = 0.75\%$, the ELM frequency increase. However, for $\leftrightarrow = 1\%$ the toroidal velocity at the top of the pedestal become counter-current and the ELM frequency decreases. The plasma density has a large influence on the range of rotation values that can be reached with changing TF ripple.

ACKNOWLEDGMENTS

This work has been conducted under the European Fusion Development Agreement (EFDA), as well as within the framework of the Contract of Association between the European Atomic Energy Community and the Instituto Superior Técnico (IST), and it has also received financial support from the Fundação para a Ciência e a Tecnologia (FCT, Lisbon). The content of the publication is the sole responsibility of the authors and does not necessarily represent the views of the European Commission, the FCT, the IST, or their services.

REFERENCES

[1]. ITER Physics Basis, *Nuclear Fusion*, **47**, S18 (2007).
[2]. T.E. Evans et al., *Physics Revie Letters*, **92**, 235003 (2004).
[3]. A.W. Degeling et al., *Plasma Physics Controlled Fusion*, **45**, 1637 (2003).
[4]. H. Urano et al., *Plasma Physics Controlled Fusion*, **45**, 1571 (2003).
[5]. Y. Sakamoto et al., *Plasma Physics Controlled Fusion*, **46**, A299 (2004).
[6]. P. de Vries et al., *Nuclear Fusion*, **48**, 035007 (2008).
[7]. H. Urano et al., *Nuclear Fusion*, **47**, 706 (2007).
[8]. Y. Andrew et al., *Review Scientific Instruments*, **77**, 10E913 (2006).
[9]. http://www.tkk.fi/Units/AES/projects/fusion/ascot.htm



Figure 1: Time traces of plasma current a)-b), magnetic field c)-d), volume averaged electron density e)-f), diamagnetic energy g)-h),outer divertor D_α i)-j) and additional heating power k) -l) for Pulse No's: 77108 and 77110.



Figure 4: (left) Angular momentum density profiles for Pulse No's: 69858 (—) and 69837 (—). (right) torque profile calculated by PENCIL, for Pulse No's: 69858 (—) and 69837 (—), and calculated by ASCOT for the latter (- - -).