

S.N. Gerasimov, T.C. Hender, J. Morris, V. Riccardo, L.E. Zakharov
and JET EFDA contributors

The Rotation of Plasma Current Asymmetries During Disruptions in JET

“This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

“Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

The contents of this preprint and all other JET EFDA Preprints and Conference Papers are available to view online free at www.iop.org/Jet. This site has full search facilities and e-mail alert options. The diagrams contained within the PDFs on this site are hyperlinked from the year 1996 onwards.

The Rotation of Plasma Current Asymmetries During Disruptions in JET

S.N. Gerasimov¹, T.C. Hender¹, J. Morris¹, V. Riccardo¹, L.E. Zakharov²
and JET EFDA contributors*

JET-EFDA, Culham Science Centre, OX14 3DB, Abingdon, UK

¹*EURATOM-CCFE Fusion Association, Culham Science Centre, OX14 3DB, Abingdon, OXON, UK*

²*Princeton Plasma Physics Laboratory, James Forrestal Campus, Princeton, NJ 08543, New Jersey, USA* See annex of F. Romanelli et al, "Overview of JET Results", (23rd IAEA Fusion Energy Conference, Daejeon, Republic of Korea (2010)).*

Preprint of Paper to be submitted for publication in Proceedings of the
39th European Physical Society Conference on Plasma Physics, Stockholm, Sweden
2nd July 2012 - 6th July 2012

INTRODUCTION

Toroidal variations in the measured plasma currents (I_p) have been observed on JET during VDEs (Vertical Displacement Events) [1],[2],[3]. The theoretical explanation of the I_p asymmetries, based on the JET disruption database, is that a long lasting $m=n=1$ kink mode is accompanied by negative helical surface plasma currents (Hiro currents) which have a pathway inside the vacuum vessel (or “machine wall”) [5], [6], see figure 1. The $m=n=1$ kink modes are responsible for the sideways forces, which occur during asymmetrical VDE disruptions. The disruptions have become a serious issue for future large-scale tokamaks. For example, the sideways forces on the vessel are expected to be tens of times greater on ITER in comparison with JET: $F_x \propto B_T I_p a$, $F_x^{ITER} \cong 2.5 \cdot 2 F_x^{JET} \cong 20 F_x^{JET}$ [3], [6].

Apart from the forces itself, the force durations (or impulse) and force time behaviour are important for the vessel structure loads. Mode frequencies that are close to the structural natural frequencies of the machine components can cause major dynamic amplifications of the loads.

PLASMA CURRENT ASYMMETRIES (HIRO CURRENT) DISRUPTION DATABASE

Replacement of carbon plasma-facing components (referred to here as JET “C-wall”) by solid beryllium limiters and beryllium tiles in the main chamber, and a combination of bulk W and W-coated divertor tiles (referred to here as JET “IL-wall”) was completed on JET in 2011 [7],[8]. The mode rotation analysis requires I_p measurements in 4 octants, each separated by 90° [3]. Only disruptions with $|I_p^{dis}| \geq 1\text{MA}$ have been analysed, where I_p^{dis} is pre-disruptive plasma current, defined as the average I_p over 20-50ms before the disruption time. The C-wall I_p asymmetries rotation database contains 951 shots. The C-wall database also contains 3483 pulses of two-octant disruption data, which has been used for non-rotational analysis. The first half year of IL-wall operation provided rotation data for 199 disruptions, and an additional 59 disruptions have only two-octant recorded data.

PLASMA CURRENT QUENCH TIME

It was known that wall material strongly effects the disruption due to impurity radiation during the Current Quench (CQ) [9]. There is a significant difference in the current decay for C-wall and IL-wall disruptions [7], [10]. The IL-wall CQ time distribution is broader and generally shifted to the range of longer decay time, figure 2. Moreover, a large fraction of IL-wall disruptions last for hundreds of milliseconds.

PLASMA CURRENT ASYMMETRIES (HIRO CURRENT) STATIC DATA

To systematically quantify the severity of I_p asymmetries the A (A_{4oct} or A_{2oct}) = $\int A_p^{asym} dt$ integral has been used, where $A_p^{asym} = I_p^{asym} / |I_p^{dis}|$, $I_p^{asym} = \sqrt{(I_{p7} - I_{p3})^2 + (I_{p5} - I_{p1})^2}$ with I_{p1} = octant 1 plasma current etc [3]. To avoid noise contributing to the results, the A integral is only evaluated for times when the start and end time window satisfied conditions: $|I_p^{asym}| > 10\text{kA}$, $A_p^{asym} > 0.5\%$,

$|I_p| > 0.1 |I_p^{dis}|$ and $|I_p^{asym}| > 20 \text{ kA}$ for the first and last 1ms window to disregard the short-lived spikes. Ignoring transients then $A \sim \int F_x dt / I_p B_t$ where F_x is the asymmetric (or sideways) force. So A is related to the magnitude of the sideways impulse force. Figure 3 shows the entire I_p asymmetry (Hiro current) data. In cases where just two orthogonal octant data values are available then a 2 octant asymmetry was defined assuming a pure sine wave in time $A = \pi A_{2oct} / 2$. The data boundary for the whole current quench duration are:

- $A = 0.10 \tau_{80-20}$ with $A_{max} = 3.7 \text{ ms}$ on the C-wall data (green lines);
- $A = 0.05 \tau_{80-20}$ with $A_{max} = 1.7 \text{ ms}$ on the IL-wall data (magenta lines).

ROTATION NUMBERS

The $m=n=1$ kink mode, which is experimentally observed on JET as I_p toroidal asymmetries, usually rotates. The mode rotation shows significant scatter in magnitude, frequency and direction [3], [6]. The 4 octant JET magnetic diagnostics allows the extraction of reliable information about toroidal mode rotation during disruptions. Fig.4 shows distinctive asymmetrical disruptions and the traces of the tip of vector $\delta I_p^*(t) = \delta I_{51}(t) e_x^* + \delta I_{73}(t) e_y^*$.

Mode rotations have sporadic behaviour. This indicates that plasma wall interactions are specifically responsible for rotation rather than due to a plasma core related effect. There is no understanding of the rotational physics or appropriate scaling at the present point in time.

Statistical analysis requires the use of criteria to extract a subset to avoid the noise contribution. Only shots satisfying $A = \int A_p^{asym} dt > 0.5 \text{ ms}$ (see Fig.3) condition have been used for the rotation statistical analysis. So the rotational statistic has been reduced from 951 to 155 shots for C-wall and from 199 to 54 shots for IL-wall. The number of rotations during disruption was defined as $N = (\varphi_{max} - \varphi_{min}) / 2\pi$. The N distributions are very similar for C- and IL- walls. However the mode rotation slightly increased for the IL-wall $\mu_N = 2.1$ ($\sigma_N = 1.2$) in comparison with the C-wall $\mu_N = 1.6$ ($\sigma_N = 0.8$), figure 5.

ROTATION FREQUENCIES

An additional condition has been applied for the frequency statistical analysis of the observed rotations. Analysis was only performed for pulses where the rotation exceeded one full turn during a disruption. As a result of this constraint, the total number of the asymmetry rotated shots was reduced to 103 shots for C-wall and to 47 shots for IL-wall.

The rotation frequency, presented in the current analysis, has been calculated as $f = 1/\tau$, where τ is the one turn period. The average $\langle A_p^{asym} \rangle$ plasma current asymmetry (Hiro current) amplitude has been calculated during

The above described algorithm has been applied to C- and IL- walls disruptions independently, see figure 6. On the next step the data points have been divided into five $\langle A_p^{asym} \rangle$ regions: $\langle A_p^{asym} \rangle < 2\%$, $2\% \leq \langle A_p^{asym} \rangle < 4\%$, $4\% \leq \langle A_p^{asym} \rangle < 6\%$, $6\% \leq \langle A_p^{asym} \rangle < 8\%$, $\langle A_p^{asym} \rangle \geq 8\%$. The means and the standard deviations of the distributions have been calculated and plotted in figure 7, where it can be seen that, at least, the mode amplitude does not decrease with frequency.

SUMMARY

1. Plasma current quench time is significantly increased for IL-wall compared with C-wall disruptions. In spite of this, the observed I_p toroidal asymmetry integral (\sim sideways force impulse) did not increase for IL-wall disruptions. It remains inside the C-wall data domain.
2. The $m=n=1$ kink mode rotation during I_p quench has sporadic behaviour. Distributions of the number of rotations are very similar for both C- and IL-wall disruptions, although rotations slightly increased for the IL-wall. Multi-turn mode rotations were observed for C- and IL-wall disruptions.
3. The Hiro current (I_p toroidal asymmetry) amplitude seems to have no degradation with mode rotation frequency for both C- and IL-walls disruption data.

Dynamic amplification remains a serious issue since high amplitude multi-turn $m=n=1$ kink mode (which are responsible for the sideways forces) rotation can cause mechanical resonance of the machine components.

ACKNOWLEDGEMENT

This work, supported by the European Communities under the contract of Association between EURATOM and CCFE, was carried out within the framework of the European Fusion Development Agreement. The views and opinions expressed herein do not necessarily reflect those of the European Commission. This work was also part-funded by the RCUK Energy Programme under grant EP/I501045.

REFERENCES

- [1]. P. Noll et al. In Proceedings of the 19th Symposium on Fusion Technology, Lisbon, 1996 , Vol. **1**, p. 751.
- [2]. V. Riccardo et al, 2010 Plasma Physics and Controlled Fusion **52** 124018.
- [3]. S.N. Gerasimov et al. P4.121, 37th EPS conference on Plasma Physics (2010), Dublin, Ireland ECA Vol. 34A.
- [4]. T.C. Hender et al. EXS/10-3 Proceedings of the 23rd IAEA FEC 2010, Daejeon, Korea.
- [5]. L. Zakharov. Physics of Plasmas **15**, 062507 (2008).
- [6]. L.E. Zakharov, S.A. Galkin, S.N. Gerasimov. Physics of Plasmas **19**, 055703 (2012).
- [7]. P. de Vries et al. I2.109, this conference.
- [8]. S. Brezinsek et al. I3.005, this conference.
- [9]. G.R. Harris. JET-R (90) 07.
- [10]. M. Lehnen et al. to be published in 24th IAEA Fusion Energy Conference (FEC2012), San Diego, USA




Figure 1: Dark blue colour represents negative Hiro current shared between vacuum vessel and plasma in VDE due to $m = n = 1$ kink mode Inter Discrete Coils are used for I_p measurements in four octants.




Figure 2: Normalised plasma current during IL-wall (left), C-wall (centre) disruptions and their CQ time distributions (right); where τ_{80-20} is the CQ time extrapolated from time to quench from 80 to 20% of I_p^{dis} .




Figure 3: The severity of I_p asymmetries (Hiro currents) for C- and IL-walls.




Figure 4: Waveforms of the the I_p , A_p^{asym} , mode toroidal angle (ϕ) and JET top view on trajectories of the tip of I_p asymmetry vector: multi-turn fast rotation (top), rotation with reversal (bottom).




Figure 5: Distribution of the number of rotations.




Figure 6: Normalised amplitude of plasma current asymmetry divided on the five $\langle A_p^{\text{asym}} \rangle$ regions versus “one turn” rotation frequency.




Figure 7: Variation of the “one turn” frequency versus average I_p asymmetry (Hiro current) amplitude.