

E.R. Solano, E.A. Autricque, I. Coffey, E. Delabie, E. de la Luna,
P. Drewelow, E. Lerche, L. Frassinetti, M. Clever, I. Nunes, T. Loarer,
A. Meigs, F. Rimini, M. Stamp, G. Sips, J. Svensson, P. Tamain
and JET EFDA contributors

Effect of Fuelling Location on Pedestal and ELMs in JET

“This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

“Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

The contents of this preprint and all other JET EFDA Preprints and Conference Papers are available to view online free at www.iop.org/Jet. This site has full search facilities and e-mail alert options. The diagrams contained within the PDFs on this site are hyperlinked from the year 1996 onwards.

Effect of Fuelling Location on Pedestal and ELMs in JET

E.R. Solano^{1,2}, E.A. Autricque³, I. Coffey⁵, E. Delabie⁴, E. de la Luna¹,
P. Drewelow⁵, E. Lerche⁶, L. Frassinetti⁷, M. Clever⁸, I. Nunes⁹, T. Loarer¹⁰,
A. Meigs⁵, F. Rimini⁵, M. Stamp⁵, G. Sips², J. Svensson¹¹, P. Tamain¹²
and JET EFDA contributors*

JET-EFDA, Culham Science Centre, OX14 3DB, Abingdon, UK

¹*For ILNF-CIEMAT, Madrid, Spain*

²*EFDA Close Support Unit, Culham, UK*

³*Centrale Lyon Eng. School, France*

⁴*Association EURATOM-FOM, Nieuwegein, The Netherlands*

⁵*CCFE, Abingdon, UK*

⁶*LPP-ERM/KMS, Brussels, Belgium*

⁷*Euratom-VR Association, Stockholm, Sweden*

⁸*FZJ, Jülich, Germany*

⁹*Associação EURATOM/IST, Lisbon, Portugal*

¹⁰*IRFM-CEA, Sant-Paul-lez-Durance, France*

¹¹*IPP- Greifswald, Germany;*

¹²*ENEA, Frascati, Italy*

* See annex of F. Romanelli et al, "Overview of JET Results",
(24th IAEA Fusion Energy Conference, San Diego, USA (2012)).

Preprint of Paper to be submitted for publication in Proceedings of the
41st EPS Conference on Plasma Physics, Berlin, Germany
23rd June 2014 – 27th June 2014

INTRODUCTION

Since the introduction of the Tungsten (W) divertor in JET, as part of the ITER-Like Wall (ILW) project, we confirm the ASDEX-Upgrade finding that low fuelling, with associated infrequent large ELMs, can cause W sputtering and entrain W into the plasma [1], eventually cooling the plasma core. To avoid core W accumulation, typically large fuelling is used at JET, producing high ELM frequencies and smaller ELMs. On the other hand, large fuelling or low pumping are known to have a detrimental effect on confinement: $H_{98} \sim 0.8$ has been typical in heavily fuelled 2MA baseline plasmas, both with C wall [2] and ILW at JET [3].

It is possible that heavy fuelling localised in the divertor area drives the plasma towards detachment and cools the X-point, thereby affecting pedestal stability, and the speed at which the separatrix recovers from the ELM. To study this we varied the poloidal fuelling location in ILW plasmas with 2MA, 2.3T, $n_e/n_{e,Greenwald} \sim 0.65$, 12–14MW of NBI heating and medium fuelling levels. Two plasma shapes were studied: one with good pumping due to strikes near the pump duct, as shown in blue in Figure 1, and a more conventional one with reduced pumping, in red in Figure 1. D_α viewing lines of inner strike, X-point and outer strike are shown for blue shape. Fuelling scans were carried out from various locations: inner strike divertor area, (iDIV, circles in next plot), upper low field side (uLFS, squares) and plasma top (Top, triangles). The divertor fuelling is toroidally distributed at source, uLFS is available at 1 toroidal location, top fuelling used 2 toroidal locations simultaneously.

Shown in Figure 2 are the ELM frequencies averaged for 1s. Empirically we observe *in this scan* that when the ELM frequencies are lower than 40Hz, sudden W influxes modify ELM frequencies and can even lead to loss of H-mode. In those cases we plot the ELM frequency 0.5s before such events. When sufficient fuelling is applied the plasma can recover from transient W events and a healthy steady state can be reached. Because the cryo-pump is located in the divertor area, divertor fuelling is less efficient. Top fuelling is most effective at increasing f_{ELM} .

Shown in Figure 3 are the plasmas that had minimum fuelling required to avoid W accumulation, fuelling location and level noted at the top of each column. Divertor fuelling was insufficient to produce regular ELMs, it typically had colder pedestals and higher radiation. The time zoom shows that in many cases we observe “negative ELMs”, drops in D_α after the ELM [3,4]. Negative ELMs are produced when a dense ($> 10^{20} \text{m}^{-3}$) cold ($T_e < 1\text{--}3\text{eV}$) plasma is being viewed, since then D_α is dominated by recombination [5]. When an ELM deposits energy in that plasma T_e rises and the recombination rate drops, leading to a reduced D_α . Later D_α may increase again as the plasma cools back down, or as n_e and T_e rise (possibly elsewhere along the line of sight) in between ELMs. Modelling D_α evolution along a line of sight is very complex. Nevertheless, we can use the observation of “negative ELMs” as an indicator of cold plasma, particularly at the X-point. In that sense we see that in iDiv fuelled plasmas the inner strike and the X-point are cold, while the outer strike is not. uLFS fuelling exhibits negative ELMs only at the inner strike. Top fuelling has mostly positive ELMs. Langmuir probes shows signs of inboard detachment, slightly more so for divertor fuelling [6].

In steady plasmas in the blue shape $H_{98} \sim 0.8\text{--}0.9$. In the low pumping red shape there is no apparent effect of puffing location: considerably colder pedestals and low confinement, $H_{98} \sim 0.75$ are observed in all cases, together with negative ELMs, even when the ELM frequencies are sufficient to control W.

In a different series of pulses, meant to study the difference between hybrid and baseline plasmas [7] we find more dynamic situations. For instance in Figure 4 we see that as power is increased inner strike and X-point ELMs change from negative to positive. At the same time pedestal temperatures and H_{98y} increase.

We find in general that at medium fuelling levels hot pedestals and good confinement are associated with a hot X-point. Whether this is related to SOL flows, as suggested in [8], is not known. Another possible interpretation is that a hotter separatrix enables the pedestal top to be hotter for a given limiting gradient. But it is also possible that a hotter pedestal produces larger energy pulses that burn through the cold plasma at the X-point and produce positive ELMs.

Returning to Figure 3 note that in the case of uLFS fuelling at 12 s there is a sudden rise in radiation, which slowly decays afterwards. It is due to a sudden W influx, captured by a spectroscopic camera as a flash of W I light (400.8nm), shown in Figure 5. Coincidentally $T_{e,\text{ped}}$ drops and $n_{e,\text{ped}}$ rises, until they recover their previous values. We interpret this as evidence of a direct effect of W influx on pedestal temperature, until the W is flushed out of the pedestal region by the ELMs, but there may also be an effect of the increase in core radiation. There are various examples such as this one in our pulses, displaying a consistent pattern.

In a different example, shown in Figure 6, we do have IR measurements at the outer strike, now in the red shape. Here the arrival of a sawtooth pulse (bringing in core impurities and fast ions) triggers a W event that increases radiation at the plasma edge (shown in the edge bolometry channels) and lowers $T_{e,\text{ped}}$. Note that the ELMs themselves are affected, their amplitude is smaller during the W and radiation event, and ELM duration is longer (it changes from 3 to 7ms in IR). Eventually the various ELMs expel W from the pedestal region (bolometry shows that some of it moves inward, some outward). When the next sawtooth cleans the core, the plasma returns to its previous state.

In conclusion: in a well pumped shape, poloidal location of fuelling can affect SOL and f_{ELM} , top fueling being most efficient at achieving stationary clean plasmas. With sufficiently low fuelling and/or strong pumping evidence of hot X-points correlates with higher $T_{e,\text{ped}}$ and higher confinement in these JET plasmas in ILW. We also show how transient W events temporarily increase P_{rad} in the pedestal, reduce $T_{e,\text{ped}}$ and modify ELMs.

This work was supported by EURATOM and carried out within the framework of the European Fusion Development Agreement. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

REFERENCES:

- [1]. R. Dux et al., Journal of Nuclear Materials **390–391** (2009) 858
- [2]. G. Saibene et al., Journal of Nuclear Materials V. **241–243**, 476 (1997)
- [3]. M.N.A. Beurskens et al., 2014 Nuclear Fusion **54** 04300
- [4]. A. Loarte et al., 1998 Nuclear Fusion **38** 331
- [5]. G.M. McCracken et al., Nuclear Fusion **38** 619 (1998)
- [6]. P. Tamain et al, 21st International Conference On Plasma Surface Interactions, O33 (2014)
- [7]. J. Mailloux, O4.127 this conference.
- [8]. M.J. Schaffer et al., 1995 Nuclear Fusion **35** 1000




Figure 1.: Blue and Red plasma shapes studied and D_α lines viewing inner strike, X-point and outer strike.




Figure 2: ELM frequency achieved for different fuelling levels and locations. Solid symbol for blue shape, open for red.




Figure 3a) for blue shape and divertor, upper LFS and top fuelling: pedestal density, temperature, D_{α} from inner strike, X-point and outer strike views, total radiation.




Figure 3b) time zoom to show negative ELMs.




Figure 4: evolution of D_{α} from negative to positive ELMs, together with change in $T_{e,ped}$.




Figure 5: shower of W sparks originating near the outer strike point, following a combined sawtooth and ELM event, responsible for triggering radiation increase and $T_{e,ped}$ drop at 12s in Pulse No: 84645, shown in Figure 4.




Figure 6: effect of W influx on pedestal.