

J. Citrin, J. Garcia, T. Görler, F. Jenko, P. Mantica, D. Told, C. Bourdelle,
D.R. Hatch, G.M.D. Hogeweyj, T. Johnson, M.J. Pueschel, M. Schneider
and JET EFDA contributors

Electromagnetic Stabilization of Tokamak Microturbulence in a High- β Regime

“This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

“Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

The contents of this preprint and all other JET EFDA Preprints and Conference Papers are available to view online free at www.iop.org/Jet. This site has full search facilities and e-mail alert options. The diagrams contained within the PDFs on this site are hyperlinked from the year 1996 onwards.

Electromagnetic Stabilization of Tokamak Microturbulence in a High- β Regime

J. Citrin^{1,2}, J. Garcia², T. Görler³, F. Jenko³, P. Mantica⁴, D. Told³, C. Bourdelle²,
D.R. Hatch⁵, G.M.D. Hogeweyj¹, T. Johnson⁶, M.J. Pueschel⁷, M. Schneider²
and JET EFDA contributors*

JET-EFDA, Culham Science Centre, OX14 3DB, Abingdon, UK

¹*FOM Institute DIFFER – Dutch Institute for Fundamental Energy Research, PO Box 1207,
3430 BE Nieuwegein, The Netherlands*

²*CEA, IRFM, F-13108 Saint Paul Lez Durance, France*

³*Max Planck Institute for Plasma Physics, Boltzmannstr. 2, 85748 Garching, Germany*

⁴*Istituto di Fisica del Plasma “P. Caldirola”, Milano, Italy*

⁵*Institute for Fusion Studies, University of Texas at Austin, Austin, Texas 78712, USA*

⁶*EES, KTH, Stockholm, Sweden*

⁷*Department of Physics, University of Wisconsin-Madison, Madison, Wisconsin 53706, USA*

** See annex of F. Romanelli et al, “Overview of JET Results”,
(24th IAEA Fusion Energy Conference, San Diego, USA (2012)).*

ABSTRACT

The impact of electromagnetic stabilization and flow shear stabilization on ITG turbulence is investigated. Analysis of a low- β JET L-mode discharge illustrates the relation between ITG stabilization, and proximity to the electromagnetic instability threshold. This threshold is reduced by suprathermal pressure gradients, highlighting the effectiveness of fast ions in ITG stabilization. Extensive linear and nonlinear gyrokinetic simulations are then carried out for the high- β JET hybrid discharge 75225, at two separate locations at inner and outer radii. It is found that at the inner radius, nonlinear electromagnetic stabilization is dominant, and is critical for achieving simulated heat fluxes in agreement with the experiment. The enhancement of this effect by suprathermal pressure also remains significant. It is also found that flow shear stabilization is not effective at the inner radii. However, at outer radii the situation is reversed. Electromagnetic stabilization is negligible while the flow shear stabilization is significant. These results constitute the high- β generalization of comparable observations found at low- β at JET. This is encouraging for the extrapolation of electromagnetic ITG stabilization to future devices. An estimation of the impact of this effect on the ITER hybrid scenario leads to a 20% fusion power improvement.

1. INTRODUCTION

The ion-temperature-gradient (ITG) instability [1, 2] is well established as a primary driver for ion and electron turbulent heat transport in tokamak plasmas [3]. While long studied, open questions remain on aspects of ITG turbulence. Specifically, experimental observations in JET tokamak L-mode discharges of significantly reduced ion temperature profile stiffness [4, 5] within the tokamak plasma half-radius – hitherto unexplained by direct numerical simulation – have recently triggered intense investigation. “Stiffness” is defined here as the gradient of the gyroBohm normalized ion heat flux with respect to the driving normalized logarithmic ion temperature gradient R/L_{Ti} . While the original experimental hypothesis posed a connection between the reduced stiffness and concomitant low magnetic shear and high $E \times B$ flow shear, this was not borne out by dedicated nonlinear gyrokinetic simulations. Rather, following an extensive numerical study using the Gene gyrokinetic code [6], the physical mechanism able to explain the observations is nonlinear electromagnetic stabilization of ITG turbulence [7, 8].

Linear stabilization of ITG modes by electromagnetic (EM) effects (i.e., finite- β) is well known [9, 10]. However, recent *nonlinear* simulations of finite- β ITG turbulence have highlighted that nonlinear EM-stabilization – i.e. the relative reduction of the heat fluxes compared with the electrostatic (ES) case – is significantly greater than the relative reduction of the linear growth rates between the EM and ES cases [11, 12]. This is correlated with an increase in zonal flow drive in the EM case [13]. When including fast ions in the system, the suprathermal pressure gradients further increase the EM coupling, augmenting the stabilization effect. This was first observed in linear calculations [14]. The nonlinear enhancement of the total EM-stabilization over the linear stabilization is maintained in the system including fast ions. In Refs. [7, 8] this was key to explain

the low T_i -stiffness observations in Refs.[4, 5]. There, the low stiffness plasmas were dominated by suprathermal pressure within halfradius. This finding has positive ramifications for extrapolation to ITER and reactor performance, where rotation is expected to be low but fusion α -particles will provide a significant source of suprathermal pressure.

Since the discharges studied in Refs. [7, 8] were relatively low- β ($\approx 0.5\%$), a natural question arises: does this physics change in more reactor-relevant, high- β plasmas? An investigation of this question is the topic of the present paper. A JET hybrid scenario, 75225 (with C-wall) [15], was chosen for detailed linear and nonlinear gyrokinetic study. The importance of fast ions on ITG stabilization in this discharge is already seen in previous linear studies [16]. In general, the enhancement of plasma diamagnetism by the suprathermal pressure gradient is believed to be essential for attaining hybrid regimes [17]. This encouraged the comparison of the ITG stabilization physics in this discharge with the aforementioned L-mode studies. Hybrid scenarios are natural candidates for such analysis due to: a significant suprathermal pressure fraction, arising from relatively low plasma density; low magnetic shear in the inner half-radius, experimentally observed to be correlated with reduced T_i stiffness [18]; and no deleterious MHD activity, which otherwise would interfere with microturbulence transport studies. The nonlinear studies shown here constitute an extension of results shown in Ref. [19], which also discusses the importance of fast ions for edge confinement improvement, leading to a beneficial core-edge coupling. The importance of fast ion enhanced linear EM-stabilization in this discharge is also predicted by the TGLF quasilinear transport model [20] when applied in an integrated modeling framework [21].

This paper concentrates on the comparison between well resolved and extensive direct numerical simulation with experimental power balance fluxes, and the ramifications thereof on relevant transport questions and extrapolations to future devices. The fundamental theory of the physics mechanisms discussed here is to a large extent still an open question, and left for future work.

The rest of the paper is organized as follows. In section 2 the simulation and discharge parameters are reviewed. Section 3 discusses the relevant parametrization of the EM-stabilization. Section 4 reviews the linear analysis of discharge 75225. Section 5 describes the nonlinear simulations of discharge 75225. Conclusions are presented in section 6.

2. SIMULATION SETUP AND DISCHARGE PARAMETERS

Gene solves the gyrokinetic Vlasov equation, coupled self-consistently to Maxwell's equations, within a δf formulation. Gene works in field line coordinates, where x is the radial coordinate, z is the coordinate along the field line, and y is the binormal coordinate. Both an analytical circular geometry model (derived in Ref.[22]) as well as a numerical geometry were used in this work. All simulations carried out were local. Collisions are modeled using a linearized Landau-Boltzmann operator. Both linear and nonlinear initial value simulations were performed. Typical grid parameters were as follows: perpendicular box sizes $[L_x, L_y] = [250, 125]$ in units of ion Larmor radii, perpendicular grid discretisations $[n_x, n_{ky}] = [256, 32]$, 32 point discretisation in the parallel direction, 48 points in

the parallel velocity direction, and 12 magnetic moments. The high L_x and n_x values were necessary to satisfy the boundary conditions in the low magnetic shear simulations at low radii, and were relaxed for simulations with higher magnetic shear. Parallel magnetic fluctuations were included in the simulations of JET discharge 75225. This cannot be neglected, due to the relatively high β , and is important for setting the strength of electromagnetic coupling, as seen in dedicated checks.

JET discharge 75225 was analyzed at two separate radial locations, at $\rho = 0.33$ and $\rho = 0.64$, where ρ is the normalized toroidal flux coordinate. The discharge is characterized by a peaked T_i profile in the inner halfradius, motivating this choice of separating the analysis to the two separate regions, and examining the relative impact of EM-stabilization and $E \times B$ flow shear stabilization in each region. For preparing the Gene input, interpretative simulations of the discharge with the CRONOS integrated modeling suite [23] were carried out. These simulations included calculations of the NBI heat deposition and fast ion profile using the NEMO/SPOT code [24], power balance analysis, current diffusion, and magnetic equilibrium calculations with HELENA [25]. The dimensionless parameters for the discharge in the two regions, included as input for the Gene simulations, are shown in Table. 1. The fast ions were approximated in Gene as hot isotropic Maxwellians, taking the average T_{fast} from the calculated slowing down distribution.

3. PARAMETRIZATION OF EM-STABILIZATION

Before proceeding to the analysis of JET discharge 75225, we first discuss the parametrization of the strength of the linear and nonlinear EM-stabilization. This is to gain intuition on the relevant parameter regimes of the effect. Since the linear stabilization of the toroidal ITG mode is closely linked with the increase of the electromagnetic instability drive [26], we take $\beta/\beta_{\text{crit}}$ as a valid parameter of merit. This will be justified in this section.

β_{crit} is the kinetic β limit of the EM ion mode, e.g., Kinetic Ballooning Modes (KBM), fast ion driven Alfvénic modes, or more generally a hybrid case. β_{crit} is a function of the plasma parameters, including the driving gradient lengths. In the fluid limit the stability threshold is parameterized by α_{MHD} , where $\alpha_{\text{MHD}} \equiv q^2 R \beta' = q^2 \sum_j \beta_j \left(\frac{R}{L_{Tj}} + \frac{R}{L_{nj}} \right)$, summing over all species. β_{crit} and α_{crit} are thus intrinsically linked, where α_{crit} effectively captures a subset of the parameter dependencies in β_{crit} . These parametrizations approximately carry over into the kinetic system [11], and thus an increase of the ITG stabilization with pressure gradients is expected (through β'). This is the key factor underlying the mechanism of fast ion stabilization of ITG, since the suprathermal pressure can increase the total pressure gradient (and thus β') and increase the stabilization factor, while not simultaneously adding to the underlying ITG drive.

A study of a subset of the β_{crit} dependencies was carried out for the low-stiffness low- β JET discharge 66404 at $\rho = 0.33$, discussed in detail in Ref.[7, 8, 27]. Of particular interest is the magnetic shear (\hat{s}) dependence, motivated by the observation of increased destiffening at low- \hat{s} [18]. In figure 1 a β scan is shown for various levels of \hat{s} , where $\hat{s} = 0.7$ is the nominal value for the discharge. These scans include both linear calculations and corresponding nonlinear simulations.

In all cases here, only the thermal species are included, although nominally 66404 has a significant fast ion population. This is an extension of the \hat{s} -scan for this discharge shown in Refs.[7, 8]. The linear runs were carried out at $k_y = 0.2$, the wavenumber corresponding to the most unstable KBM mode. k_y is normalized by $1/\rho_s$

For all values of \hat{s} , a transition from ITG modes to KBM is observed, characterized by a sharp upswing of the growth rates for increasing β and a jump in the real frequencies of the dominant mode. Positive ω is defined as the ion diamagnetic direction in Gene. For $\hat{s} > 1$, an intermediate TEM zone is apparent, characterized by the negative frequencies. The KBM thresholds for each case were extrapolated by a quadratic fit to the first several KBM growth rates, shown in figure 1a.

An \hat{s} -scan of nonlinear simulations – both EM and ES – was then carried out, with the nominal $\beta_e = 0.32\%$ corresponding to discharge 66404 at $\rho = 0.33$. The ratio between the ES-case and EM-case ion heat fluxes at each point in the scan is shown in figure 1c, plotted against the $\beta_e/\beta_{\text{crit}}$ for each parameter set, where β_{crit} was taken from the fit of the linear scans. Two important points should be emphasized: a) the β scans and β_{crit} shown here correspond to electron β , but the β_s of all ionic species in the simulations are scaled self-consistently with β_e , according to the pressure ratios p_j/p_e ; b) the \hat{s} -dependent β_{crit} is the parameter which changes within the scan in figure 1c, not β_e . While the intermittency in the simulations leads to significant error bars, a clear separation in this flux ratio is seen between high and low $\beta_e/\beta_{\text{crit}}$, with a higher ratio at higher $\beta_e/\beta_{\text{crit}}$. This signifies that $\beta/\beta_{\text{crit}}$ is a valid parameter of merit also for the nonlinear EM-stabilization, suggesting that the EM modification of the linear modes is also linked to the enhanced nonlinear stabilization.

In figure 2, the relevance of the $\beta/\beta_{\text{crit}}$ parametrization of EM-stabilization in the presence of fast ions is examined. Figure 2a displays a β -scan for three separate cases of discharge 66404 at $\rho = 0.33$: with thermal species only, with a fast D species matching the modeled NBI profile for the discharge, and with a fast ^3He species matching the modeled ICRH profile. In this scan, a separation of the curves are evident, with the EM-stabilization occurring at different rates with increasing β_e . Continuing the calculation of the curves down to the KBM limit (not shown for brevity) we obtain $\beta_{\text{crit}} = 0.77\%$, 0.66% , 0.59% respectively for the three cases. When rescaling the xaxis of the plot to $\beta_e/\beta_{\text{crit}}$, as shown in figure 2b, the overlap between the three curves is much more apparent, illustrating the suitability of $\beta/\beta_{\text{crit}}$ as a parameter of merit.

In conclusion, $\beta/\beta_{\text{crit}}$ is shown to be a valid parameter of merit for both the linear and nonlinear EMstabilization. β_{crit} is reduced at lower- \hat{s} and with increased suprathermal pressure gradients. Thus, for a given thermal β , enhanced EM-stabilization of ITG turbulence is expected at both low- \hat{s} and high suprathermal pressure. This statement is relevant for the interpretation of transport in JET discharge 75225, as shown in the following sections.

4. LINEAR SIMULATIONS OF JET 75225

In this section we describe the linear analysis of the JET high- β hybrid discharge 75225. The analysis is split into 2 subsections concentrating on different radii, $\rho = 0.33$ and $\rho = 0.64$.

4.1. LINEAR CALCULATIONS AT $\rho = 0.33$

In figure 3 we plot the k_y spectrum of the linear growth rates and frequencies at $\rho = 0.33$. The plot shows 4 separate cases: a case without fast ions, the nominal case including the full fast ion pressure gradient, a case with the fast ion pressure gradient reduced by 30%, and a case where β_e was set to 10^{-5} , which is effectively an electrostatic case. The spectrum in all cases is dominated by ITG modes, characterized by the positive frequencies in the diamagnetic drift frequency range. From the comparison of the EM and ES cases, a strong linear EMstabilization is evident, particularly at the higher k_y values.

An additional mode is clearly apparent for the nominal parameters, at $k_y < 0.2$, at significantly higher frequencies than the ITG range. This mode is stabilized by lowering the fast ion pressure gradient by 30%, illustrating the fast ion drive of this mode. From a Rosenbluth-Hinton test [28] it was determined that the frequency of these modes are within 5% of the GAM frequency for these parameters. This supports the identification of this mode as a Beta induced Alfvén Eigenmode (BAE) [29], which is known to be degenerate with the GAM frequency [30]. However, these modes are likely coupled with the KBM modes which share a similar frequency range. Thus, in lieu of deeper analysis regarding the precise characterization, in the following we refer to this mode as a hybrid BAE/KBM. As will be shown in section 5, the heat fluxes in the simulation of the system with unstable BAE/KBMs is highly inconsistent with the experimental power balance values. Thus, it is assumed that the system selforganizes to a state just below the BAE/KBM stability boundary, and we maintain the reduced fast ion pressure for all the nonlinear simulations.

Comparing the ITG growth rates of the cases with and without fast ions shows that the fast ions provide an additional stabilization of ITG beyond the thermal EM-stabilization, as expected from section 3. This stabilization is only slightly reduced when lowering the fast ion pressure by 30%. The magnetic equilibrium modified by the fast ions is self-consistently included in the calculation, and therefore the increased stabilization when including fast ions includes a contribution from both EM and geometric (Shafranov shift) stabilization. However, from dedicated checks, the EM stabilization dominates in this case. However, we note that the Shafranov shift stabilization is critical for improving peeling-ballooning stability in the tokamak edge region, meaning that fast ions can improve pedestal stability, leading to a beneficial core-edge transport coupling [19].

The behavior of the BAE/KBM mode destabilization at $k_y = 0.1$ is shown in figure 4, for the two separate cases with nominal and reduced fast ion pressure gradient. For both cases the ITG growth rates are stabilized with higher β_e until the BAE/KBM is eventually destabilized. For the nominal gradients, this destabilization occurs for a β_e lower than the experimental β_e of 1.84%. For the reduced fast ion pressure gradient, the instability boundary is just above the experimental value. Thus, $\beta/\beta_{\text{crit}} \approx 1$ in this case, and according to the results in section 3 we can thus expect significant nonlinear EM-stabilization as well. We reiterate that in the scan, the β_s of all ionic species are also scaled in the scan with β_e , according to the pressure ratios p_j/p_e .

4.2. LINEAR CALCULATIONS AT $\rho = 0.64$

In figure 5 we show the linear calculations at $\rho = 0.64$. Three cases are shown: the nominal case including fast ions, a case without fast ions, and an electrostatic case. Three distinct instabilities are apparent. In the EM cases, a micro-tearing-mode (MTM) – identified from its mode parity and structure – is unstable at $k_y = 0.05$. For intermediate k_y ITGs are unstable, and for $k_y > 1.1$ trapped electron modes (TEMs) are unstable. As opposed to the $\rho = 0.33$ case, the EM-stabilization is much weaker in the lower, transport driving k_y values. This is reflected by the relatively lower $\beta/\beta_{\text{crit}} \approx 0.3$ values calculated for this case.

We note that these linear calculations were carried out for modified input parameters compared with table 1, with R/L_{Te} and R/L_{ne} reduced by 20%, and R/L_{Ti} increased by 20%. This was necessary to reach agreement with the experimental power balance values and q_i/q_e heat flux ratio in the nonlinear simulations. For nominal parameters, a strongly driven TEM regime was predicted at all k_y , leading to $q_e > q_i$, in disagreement with the observations. This modification of the driving gradients is at the limits of the experimental error bars.

5. NONLINEAR SIMULATIONS OF JET 75225

In this section we describe the nonlinear simulations of JET discharge 75225, split into 2 subsections concentrating on different radii, $\rho = 0.33$ and $\rho = 0.64$.

5.1. NONLINEAR SIMULATIONS AT $\rho = 0.33$

In figure 6 the Gene saturated nonlinear transport fluxes are shown for various cases: with fast ions (with 30% reduced suprathermal pressure gradient), without fast ions, and with fast ions but electrostatic. In addition, the experimental ion and electron heat fluxes from power balance is shown. The Gene error bars are derived from intermittency and uncertainties in the precise saturated potential level. The power balance ion heat flux uncertainties include statistical errors, and the propagation of $T_{i,e}$ errors on the ion-electron heat exchange term.

The most striking observation is that the nonlinear EM-stabilization is absolutely critical for achieving agreement between the Gene and experimental flux values. The nonlinear EM-stabilization is here manifested mostly as a stiffness reduction. The approximate factor 20 reduction in heat flux is much greater than the analogous growth rate reduction, reflecting the enhanced nonlinear nature of the EM-stabilization. This result is thus a reactor-relevant high- β extension of the results reported in Ref. [5, 7]. In addition to the stiffness reduction, a slight increase in the value of the Dimits shift [31] is observed between the ES and EM case with fast ions. This is seen by comparing the linear R/L_{Ti} thresholds for these cases, calculated separately and shown in the figure, with the extrapolation to zero-flux of the nonlinear stiffness curves. This Dimits shift extension is consistent with previous findings of nonlinear EM-stabilization [11, 12]. In this case, as opposed to the low- β cases in Ref. [7], the majority of the nonlinear EM-stabilization is due to the thermal component. This is consistent with the lower suprathermal pressure fraction in these high- β cases compared with the low- β cases,

on the order of 30% here. However, the addition of fast ions does still provide significant additional stabilization, leading to an additional increase of R/L_{Ti} by approximately 10–20%, for the same heat flux level.

The level of agreement between the experimental power balance and Gene predictions is remarkable, and provides a highly encouraging validation of nonlinear gyrokinetics in the high- β regime. This is only possible due to the increasing capacity in high performance supercomputing capabilities. For example, the simulations shown in figure 6 necessitated around 4 million CPU hours of calculation time, not including extensive convergence tests. The rightmost datapoint in figure 6, with its sharp uptick in heat fluxes at the highest R/L_{Ti} , is a consequence of the destabilization of the BAE/KBM mode, which is also destabilized by the main ion gradients. This is also seen by the strong increase in fast ion diffusivity for that case. The particularly sharp uptick of the electron heat flux is mostly due to increased magnetic flutter. We recall that these simulations all used a suprathermal pressure gradient reduced by 30%. The anomalous diffusivity prescribed in the SPOT code to provide such a suprathermal pressure reduction in the CRONOS interpretative simulations was $0.5m^2/s$. This level is actually consistent with the Gene predicted fast ion diffusivity for the simulation at nominal R/L_{Ti} , which also led to heat flux agreement with power balance. Thus, the selfconsistency of this assumption is verified.

In ITG turbulence, zonal flows (ZFs) are a well known saturation mechanism [32]. The correlation of the nonlinear EM-stabilization and increased destabilization of ZF secondaries has been reported [33]. This connection is verified in these simulations as well. This is seen by analyzing the ratio of the amplitude of the ZFs (all $k_y = 0$ modes) to the sum of amplitudes of all drift wave modes. This value, $|\phi_{ZF}^2| / \sum_{k_y > 0} |\phi_{DW}^2|$, serves a proxy for the relative impact of ZFs in the nonlinear system. This ratio is ≈ 40 for the EM case with fast ions, ≈ 30 for the EM case without fast ions, and ≈ 10 for the ES case. This trend correlates well with the trend of the relative heat flux ratios. In figure 7, the normalized nonlinear amplitude spectra is shown for the nominal EM case and ES case. The relative increased strength of the $k_y = 0$ ZF modes in the EM case is clearly evident in the plot.

The relative role of EM-stabilization and flow shear stabilization was investigated by γ_E scans. This is shown in figure 8. The γ_E scans, where $\gamma_E = 0.2$ is the nominal value, were carried out for both EM and ES cases. A number of striking observations come out from this scan. Firstly, for the (nominal) EM case, no flow shear stabilization was evident at all. A slight increase in fluxes is even observed, particularly in the electron channel. This increased q_e is due to increased magnetic flutter, which may suggest increased fast ion mode destabilization with rotation. For the ES case, a more classical trend is observed, with stabilization with increased γ_E . However, for the nominal $\gamma_E = 0.2$, this stabilization remains weak. Even with $\gamma_E = 0.6$, the flux levels are significantly above the power balance levels. This suggests that rotation is not an important stabilization factor in this regime, which is dominated by EM-stabilization, at least for inner radii such as $\rho = 0.33$. The explanation for the complete lack of flow shear stabilization in the EM case remains unknown, and will be studied in the future. A conjecture is that this may be related

to the much shorter autocorrelation times of the turbulent structures in the EM cases compared with the ES cases.

We conclude this subsection with a brief discussion of the combined ITG-BAE/KBM turbulent state. This was carried out by carrying out a simulation using the nominal fast pressure gradient, i.e. the case in which the BAE/KBM mode is seen to be unstable in figure 3. The time dependent nonlinear fluxes are shown in figure 9. In the plot, an initial phase is evident, corresponding to the case with reduced fast pressure gradient. The simulation input is then shifted to the nominal fast pressure gradient, and the simulation is restarted. This then corresponds to phase 2, where all fluxes rise to a degree significantly above the power balance levels. In particular, the EM component (magnetic flutter) of the electron heat flux rises sharply. This EM transport component is correlated with tearing parity fluctuations at $k_y > 0.3$. MTMs are linearly stable at this region, and this observation is potentially linked to heightened nonlinear mode-mixing to tearing parity modes [34] when the BAE/KBM is unstable. This is a subject of future study. However, this regime as an operating point is not experimentally relevant due to the high fluxes, and this supports the use of “stiff” Alfvén Eigenmode models in reduced modeling frameworks [35]. Whether the system is maintained constantly below the BAE/KBM mode limit, or has limit cycle dynamics, is an open question and necessitates flux driven simulations. Thus, we cannot fully rule out the experimental relevance of this state. Finally, we note that the phase 1 time trace shown in figure 9 is the tail end of a significantly longer simulation, with sufficient statistics to determine a saturated nonlinear state. However, for phase 2, there are insufficient statistics for such a determination. Nonetheless, we deem the qualitative observation of significantly higher fluxes than the power balance as robust.

5.2. NONLINEAR SIMULATIONS AT $\rho = 0.64$

In this subsection we describe the nonlinear analysis at $\rho = 0.64$, and compare the trends observed with the analogous simulations at $\rho = 0.33$. The R/L_{T_i} scans are shown in figure 10. Numerous scans were carried out, isolating the relative roles of EM-stabilization (thermal and with fast ions), and $E \times B$ flow shear stabilization. It is clear that at this location, the EM-stabilization is extremely weak, with the ES and EM curves in close proximity. This is likely due to the much lower $\beta/\beta_{\text{crit}} \equiv 0.3$ in this case compared with $\rho = 0.33$.

The main observation is that flow shear stabilization is effective for these parameters, and can significantly increase the R/L_{T_i} values which match the power balance fluxes. In general, the stiffness level is much higher for these parameters compared with $\rho = 0.33$, likely due to the higher \hat{s} and q values as well as lower EMstabilization. The $E \times B$ stabilization provides a threshold shift.

We reiterate that these nonlinear simulations were carried out for modified input parameters compared to the nominal parameters from table 1, with R/L_{T_e} and R/L_{ne} reduced by 20%, and R/L_{T_i} increased by 20%. This was to avoid a strongly driven TEM regime and match the experimental q_i/q_e , and reach the vicinity of the absolute flux values.

5.3. EXTRAPOLATION TO ITER

We conclude this section with a rough extrapolation of the inner core EM-stabilization effect to ITER, with the aim of judging the scale of potential impact. Theory based extrapolations carried out until now have not included the enhanced nonlinear EM-stabilization effects. This extrapolation was based on a T_i profile predicted for an ITER hybrid scenario in previous theory based integrated modeling [36]. This modeling consisted of a CRONOS predictive simulation with the GLF23 transport model [37, 38], and a pedestal height of 4 keV. As seen in figure 11, this reference T_i is compared with an estimate – based on the same profile – of T_i peaking in the inner core between $\rho = 0.2 - 0.5$. This peaking was estimated by a simple $\Delta R/L_{T_i} = 2$ upshift, based on the results of JET 75225, which shares a similar β and suprathermal pressure fraction as predicted for this ITER hybrid scenario reference case. The lower cutoff point of $\rho = 0.2$ is based on a conservative assumption that access to the ITG EM-stabilization zone may be hindered by extremely low- \hat{s} in the very inner core which decreases the β_{crit} of the BAE/KBMs [39]. Based on the $n_{D,T}$ from the reference case, the fusion power corresponding to the two separate T_i profiles can be compared. The reference $P_{\text{fus}} = 350\text{MW}$ increases by 20% to $P_{\text{fus}} = 420\text{MW}$. The impact of core EM-stabilization is thus estimated to be significant.

CONCLUSIONS

Detailed linear and nonlinear local gyrokinetic calculations of the JET high- β hybrid discharge 75225 at $\rho = 0.33$ and $\rho = 0.64$ were carried out. Good agreement between the simulated and power balance heat fluxes was achieved. It was found that at $\rho = 0.33$, electromagnetic (EM) stabilization of ITG turbulence is critical for reaching agreement with power balance heat fluxes, while $E \times B$ flow shear stabilization was not effective. Nonlinear effects and the inclusion of suprathermal pressure both significantly enhance the EM-stabilization. At $\rho = 0.64$, on the other hand, the EM-stabilization was ineffective while the $E \times B$ flow shear stabilization was important. A likely explanation for the difference in EM-stabilization effectiveness at the two locations is the value of the $\beta/\beta_{\text{crit}}$ EM-stabilization parameter of merit, which is ≈ 1 and 0.3 respectively for the two cases. β_{crit} is the critical β value for the onset of EM instabilities.

These results provide a high- β extension of previous results of enhanced nonlinear EM-stabilization at low- β . The reactor relevance of this particular discharge is encouraging for extrapolations to future devices, which will have low rotation but a significant suprathermal pressure fraction due to fusion α -particles. An estimate of the impact of this effect on ITER extrapolation was carried out, with a 20% increase of fusion power predicted for an ITER hybrid scenario. This is a strong indication that reduced models employed for transport extrapolation to ITER and reactors must include suprathermal pressure and EM-stabilization effects validated by nonlinear gyrokinetic simulations.

Future work will concentrate on the physical mechanism of the enhanced nonlinear EM-stabilization, which is likely related to either (or both) increased secondary ZF growth rates or reduced tertiary damping of ZFs. However, a precise characterization of the effect is still lacking.

ACKNOWLEDGMENTS

This project has received funding from the European Unions Horizon 2020 research and innovation programme under grant agreement number 210130335, and financial support from NWO. The views and opinions expressed herein do not necessarily reflect those of the European Commission. This work is supported by NWO-RFBR Centre-of-Excellence on Fusion Physics and Technology (Grant nr. 047.018.002). This work is part of the research programme ‘Fellowships for Young Energy Scientists’ (YES!) of the Foundation for Fundamental Research on Matter (FOM), which is financially supported by the Netherlands Organisation for Scientific Research (NWO). The authors would like to thank E. Highcock, S. Moradi, G. Plunk, A. Schekochihin and E. Westerhof for stimulating discussions. Resources of the HELIOS supercomputer at IFERC-CSC are acknowledged. This research used computational resources at the National Research Scientific Computing Center, which is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. The authors are grateful to D. R. Mikkelsen for assistance.

REFERENCES

- [1]. F. Romanelli, 1989 *Physics of Fluids B* **1** 1018.
- [2]. S.C. Guo and F. Romanelli, 1993 *Physics of Fluids B* **5** 520.
- [3]. E.J. Doyle et al. Progress in the ITER Physics Basis Chapter 2: Plasma confinement and transport 2007 *Nuclear Fusion* **47** S18.
- [4]. P. Mantica et al. 2009 *Physical Review Letters* **102** 175002.
- [5]. P. Mantica et al. 2011 *Physical Review Letters* **107** 135004.
- [6]. F. Jenko, W. Dorland, M. Kotschenreuther and B.N. Rogers 2000 *Physics of Plasmas* **7** 1904, and <http://gene.rzg.mpg.de>.
- [7]. J. Citrin et al. 2013 *Physical Review Letters* **111** 155001.
- [8]. J. Citrin et al. 2014 *Nuclear Fusion* **54** 023008.
- [9]. J. Weiland and A. Hirose 1992 *Nuclear Fusion* **32** 151.
- [10]. A. Hirose 2000 *Physics of Plasmas* **7** 433.
- [11]. M.J. Pueschel, M. Kammerer and F. Jenko 2008 *Physics of Plasmas* **15** 102310.
- [12]. M.J. Pueschel and F. Jenko 2010 *Physics of Plasmas* **17** 062307.
- [13]. M.J. Pueschel, T. Görler, F. Jenko, D.R. Hatch, and A.J. Cianciara 2013 *Physics of Plasmas* **20** 102308.
- [14]. M. Romanelli, A. Zocco, F. Crisanti and JET-EFDA Contributors, 2010 *Plasma Physics and Controlled Fusion* **52** 045007.
- [15]. J. Hobirk et al., *Plasma Physics and Controlled Fusion* **54**, 095001 (2012).
- [16]. J. Garcia et al., *Nuclear Fusion* **53**, 043023 (2013).
- [17]. J. Garcia and G. Giruzzi, *Physical Review Letters* **104**, 205003 (2010).
- [18]. P. Mantica et al. 2011 *Plasma Physics and Controlled Fusion* **53** 124033.
- [19]. J. Garcia et al., submitted to *Physical Review Letters*

- [20]. G.M. Staebler et al., 2005 Physics of Plasmas **12**, 102508.
- [21]. B. Baiocchi et al., 2014 Submitted to Plasma Physics and Controlled Fusion.
- [22]. X. Lapillonne et al. 2009 Physics of Plasmas **16** 032308.
- [23]. J.F. Artaud et al. 2010 Nuclear Fusion **50** 043001.
- [24]. M. Schneider et al. 2011 Nuclear Fusion **51** 063019.
- [25]. G.T.A. Huysmans, J.P. Goedbloed and W. Kerner 1991 CP90 Conf. on Comp. Physics (Singapore: Word Scientific) p 371.
- [26]. J.Y. Kim, W. Horton and J.Q. Dong, Physics of Fluids B **5**, 4030 (1993).
- [27]. F. Ryter et al., Nuclear Fusion **51**, 113016 (2011).
- [28]. M.N. Rosenbluth and F.L. Hinton, Physical Review Letters **80**, 724 (1998).
- [29]. F. Zonca et al., Plasma Physics and Controlled Fusion **38**, 2011 (1996).
- [30]. C. Nguyen, X. Garbet and A.I. Smolyakov, Physics of Plasmas **15** 112502 (2008).
- [31]. A.M. Dimits et al. 2000 Physics of Plasmas **7** 969.
- [32]. P. Diamond, S-I. Itoh, K.Itoh and T.S. Hahm, 2005 Plasma Physics and Controlled Fusion **47** R31.
- [33]. M.J. Pueschel et al. 2013 Physical Review Letters **110** 155005.
- [34]. D.R. Hatch et al. 2012 Physical Review Letters **108** 235002.
- [35]. E.M. Bass and R.E. Waltz 2010 Physics of Plasmas **17**, 112319.
- [36]. J. Citrin et al., 2010 Nuclear Fusion **50** 115007.
- [37]. R.E. Waltz, G.M. Staebler, W. Dorland and G.W. Hammett 1997 Physics of Plasmas **4** 2482.
- [38]. Kinsey J.E., Staebler G.M. and Waltz R.E. 2005 Physics of Plasmas **47** 052503.
- [39]. S. Moradi et al., 2014 Submitted to Nuclear Fusion.

Location	R/L_{Ti}	R/L_{Te}	R/L_{ne}	T_i/T_e	β_e [%]	\hat{s}	q	ν^*	Z_{eff}	γ_E	n_{fast}/n_e	T_{fast}/T_e	$R/L_{T\text{fast}}$	$R/L_{n\text{fast}}$
$\rho = 0.33$	6.2	3.4	2.4	1.2	1.8	0.16	1.1	0.023	1.55	0.2	0.12	7.3	0.33	7.0
$\rho = 0.64$	3.4	5.8	3.05	1.07	0.75	1.44	1.74	0.018	1.90	0.3	0.055	8.65	2.15	8.9

Table 1: Discharge 75225 dimensionless parameters as input into the GENE simulations. The values are averaged between 6.0-6.5 s, before the onset of a deleterious $n = 3$ neoclassical tearing mode. ν^* is the normalized collisionality: $\nu^* \equiv \nu_{ei} \frac{qR}{\epsilon^{1.5} v_{te}}$, with $\epsilon = a/R$ and $v_{te} \equiv \sqrt{T_e/m_e}$. γ_E is the $E \times B$ shear rate, where $\gamma_E \equiv \frac{r}{q} \frac{d\Omega}{dr} / (\frac{v_{th}}{R})$ for the purely toroidal rotation assumed here for the NBI driven cases. $v_{th} \equiv \sqrt{T_e/m_i}$. The gradient lengths are defined taking the radial coordinate as the toroidal flux coordinate. We note that the β_j of all ionic species are scaled with β_e , according to the pressure ratios p_j/p_e .




Figure 1: β scans for various \hat{s} values for the low-stiffness JET Pulse No: 66404. The nominal β_e and \hat{s} is 0.32% and 0.7 respectively. The left panel (a) shows the linear growth rates and the quadratic fit to the β_{crit} KBM threshold (the curves meeting the x-axis). The center panel (b) shows the corresponding frequencies, and the right panel (c) shows the ratio between the ES-case and EM-case ion heat fluxes from the corresponding nonlinear simulations.




Figure 2: Linear β scans for JET Pulse No: 66404 at $p = 0.33$, for thermal species only and when including either NBI or ICRH driven fast ions. The scans are plotted as a function of β_e (a) and as a function of β_e/β_{crit} (b), where β_{crit} was separately calculated for each case.




Figure 3: Linear growth rate (a) and frequency (b) spectra as a function of k_y , for JET Pulse No: 75225 at $\rho = 0.33$. The calculations included numerical geometry, collisions, C impurities, EM fluctuations and fast ions, unless otherwise noted.




Figure 4: β -scan of linear growth rate at $k_y = 0.1$ for JET Pulse No: 75225 at $\rho = 0.33$. Two separate cases are shown, one with nominal parameters and the second with the fast ion pressure reduced by 30%. The nominal $\beta_e = 1.84\%$ is signified by the vertical black line.




Figure 5: Linear growth rate (a) and frequency (b) spectra as a function of k_y , for JET Pulse No: 5225 at $\rho = 0.64$. The calculations included numerical geometry, collisions, C impurities, EM fluctuations and fast ions, unless otherwise noted.




Figure 6: Ion heat fluxes (a), electron heat fluxes (b), and fast ion particle diffusivity (c) from Gene nonlinear simulations of JET Pulse No: 75225 at $\rho = 0.33$.




Figure 7: Amplitude of the electrostatic potential averaged over the saturated nonlinear phase for two separate cases (EM and ES) of JET Pulse No: 75225 at $\rho = 0.33$. The amplitudes at each k_y are averaged for all k_x modes and over the parallel direction.




Figure 8: γ_E scans for JET Pulse No: 75225 at $\rho = 0.33$, for both EM (nominal) and assumed ES cases. The nominal γ_E is 0.2. The simulations were carried out for $R/L_{Ti} = 6$, and the ion heat fluxes are compared against the power balance value.




Figure 9: Time traces of Gene simulated ion heat flux (a), electron heat flux (b), and fast ion diffusivity (c) for JET Pulse No: 75225 at $\rho = 0.33$ during a phase with a 30% reduced fast ion pressure gradient (phase 1), followed by a transition back to the nominal fast ion pressure gradient (phase 2). The time traces are compared with the power balance heat fluxes, whose 1σ confidence region is shown by the grey shaded zone.




Figure 10: Ion heat fluxes from Gene nonlinear simulations of JET Pulse No: 75225 at $\rho = 0.64$, for various cases including EM and ES simulations, and without $E \times B$ flow shear.




Figure 11: Ion temperature profiles from a reference CRONOS/GLF23 ITER hybrid scenario simulation, compared with an estimate of a peaked T_i profile when including core EMstabilization effects. The fusion powers of the 2 cases are compared.