

JET-P(89)31

B. Denne, G. Magyar, J. Jacquinot
and JET Team

Beryllium Like MO XXXIX and Lithium Like MO XL Observed in the JET Tokamak

“This document contains JET information in a form not yet suitable for publication. The report has been prepared primarily for discussion and information within the JET Project and the Associations. It must not be quoted in publications or in Abstract Journals. External distribution requires approval from the Publications Officer, JET Joint Undertaking, Abingdon, Oxon, OX14 3EA, UK”.

“Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

The contents of this preprint and all other JET EFDA Preprints and Conference Papers are available to view online free at www.iop.org/Jet. This site has full search facilities and e-mail alert options. The diagrams contained within the PDFs on this site are hyperlinked from the year 1996 onwards.

Beryllium Like MO XXXIX and Lithium Like MO XL Observed in the JET Tokamak

B. Denne, G. Magyar, J. Jacquinet
and JET Team*

JET-Joint Undertaking, Culham Science Centre, OX14 3DB, Abingdon, UK

** See Appendix 1*

Preprint of Paper to be submitted for publication in
Physical Review A

Berylliumlike Mo XXXIX and Lithiumlike Mo XL

Observed in the JET Tokamak

B. Denne, G. Magyar and J. Jacquinet

JET Joint Undertaking, Abingdon, OXON. OX14 3EA, U.K.

ABSTRACT

The resonance lines of berylliumlike and lithiumlike molybdenum have been observed in high-temperature JET tokamak discharges into which molybdenum was introduced using the laser blow-off method. The measured transition wavelengths are $2s^2\ ^1S_0 - 2s2p\ ^1P_1$ $49.904 \pm 0.03\ \text{\AA}$, $2s^2\ ^1S_0 - 2s2p\ ^3P_1$ $137.787 \pm 0.03\ \text{\AA}$ for Mo XXXIX, and $2s\ ^2S_{1/2} - 2p\ ^2P_{1/2}$ $143.998 \pm 0.02\ \text{\AA}$, $2s\ ^2S_{1/2} - 2p\ ^2P_{3/2}$ $58.499 \pm 0.02\ \text{\AA}$ for Mo XL. For the berylliumlike ion the results are compared with the semiempirical predictions by Edlén, while the results for lithiumlike Mo are compared with the calculations by Johnson *et al.*

I. INTRODUCTION

The study of highly ionized, high- Z elements is of interest for spectroscopic diagnostics of high-temperature plasmas such as are found in modern tokamaks like the Joint European Torus (JET) and the Tokamak Fusion Test Reactor (TFTR). Molybdenum ($Z=42$) in its Be- and Li-like ionization stages is suitable for diagnosing plasmas with central electron temperatures in the range 6 - 10 keV.

Further, accurate determination of the transition wavelengths for the resonance lines of few-electron systems such as the Be- and Li-like ions of high- Z elements is of fundamental interest in atomic physics, where such measurements can be compared to precise theoretical calculations or semi-empirical predictions. Thus Johnson *et al.*¹ have pointed out that for the Li-sequence, there is a small but significant discrepancy between experimental values and calculations including 1-electron QED contributions, which they have attributed to screening of the Lamb shift. For the Be-sequence resonance lines, Denne and Hinnov² have noted a systematic deviation from the semi-empirical predictions by Edlén³ in the range $Z = 28 - 36$: the singlet line wavelengths become increasingly longer and the intercombination line wavelengths shorter than predicted with increasing Z . Tokamak plasmas with high carbon concentrations (as is the case for both JET and TFTR) provide good possibilities for accurate measurements since the carbon lines emitted are strong and high grating orders of the C V and C VI resonance lines are easily observable, and may thus be used as references.

II. EXPERIMENT

The target plasmas were 3 MA and 5 MA deuterium discharges into which molybdenum was introduced using the recently installed JET laser blow-off system⁴. In addition to the ohmic heating, several MW of ion cyclotron resonance heating (ICRH), in the hydrogen minority scheme, were admitted for supplementary heating. The central electron temper-

ature was in the range 9.2 - 10.4 keV, and the central electron density ranged from 2.8 to $3.7 \times 10^{13} \text{ cm}^{-3}$ for the 3 MA discharges, whereas the corresponding parameters for the single 5 MA discharge used were 7.6 keV and $6.3 \times 10^{13} \text{ cm}^{-3}$, respectively.

Spectra were recorded using a 2-m extreme grazing-incidence Schwob-Fraenkel spectrometer⁵, equipped with a 600 g/mm grating, and two microchannel-plate image intensifier/converter detector systems, fiber-optically coupled to 1024-element photodiode arrays. The detectors, moveable along the Rowland circle, cover 20 - 60 Å portions of spectrum, depending on wavelength. The nominal spectral range of the instrument is (approximately) 15 - 340 Å. In the present experiment, however, scans were restricted to regions covering the four resonance lines, and in first grating-order only, due to a lack of machine time.

In Fig. 1 is shown the $\sim 49 - 59 \text{ Å}$ region before (dashed line) and after (solid line) molybdenum injection. The Mo XXXIX $2s^2 \ ^1S_0 - 2s2p \ ^1P_1$ and Mo XL $2s \ ^2S_{1/2} - 2p \ ^2P_{3/2}$ transitions are seen along with part of the C VI Lyman series in second grating order. These carbon lines were used as wavelength references, along with C VI L_α in second order (at 67.4720 Å) and the C V resonance and intercombination lines at 40.2683 Å⁶ and 40.7306 Å, respectively. For the two long-wavelength molybdenum lines (around $\sim 140 \text{ Å}$), again C V and C VI lines (in high orders) were used as references, as well as nickel lines (nickel being an intrinsic impurity in JET). For the wavelength determination, spectral data were integrated for 164 msec, while, in order to obtain the temporal evolution of the molybdenum lines with adequate time resolution, scans were made with 16 msec integration time. Figure 2 shows the time histories of the Mo XXXII 127.868-Å line (Na-like) and the Li-like Mo XL line at 143.998 Å. Both lines have been normalized to their peak values; on an absolute scale, the Mo XL signal is about 20 times smaller than the Mo XXXII signal - hence the much higher level of noise in the former signal. The molybdenum injection takes place at 7.5 sec into the discharge, as indicated in the Figure. During the time depicted in Fig. 2, the electron density remains constant. The central electron temperature, also shown in Fig. 2, is on the average constant, but exhibits sawtooth-like behavior, caused by a

magnetohydrodynamic instability of the plasma. It is seen in Fig. 2 that the Mo XL-line peaks considerably later (~ 250 msec) than the Mo XXXII-line, confirming the behavior expected of a much higher ionization stage. (The immediate early peak just after 7.5 sec may be attributed to a blending line of a low ionization stage of Mo, or a momentary change in the background plasma, associated with the injection.) The drop to a lower level of the Mo XL signal at ~ 8 sec is presumably caused by the sawtooth crash in the temperature at this time, leading to an expulsion of Mo from the plasma center. A corresponding increase in signal for the Mo XXXII-line may be seen at the same time, this ion being located radially further out in the plasma (where the temperature is essentially unchanged). The Mo XL signal recovers to a lower level after the crash, and a further drop can be seen at the time of the next sawtooth crash, at ~ 8.4 sec, although less pronounced.

For the wavelength determination, the positions of the molybdenum lines and the reference lines were established. A weighted, least-squares fit of the references was made using orthogonal (Chebyshev) polynomials. Generally, a fourth-order polynomial provided the best fit. The wavelengths of the Mo lines were subsequently determined.

As mentioned earlier, measurements were performed in first grating-order only. The two long-wavelength lines could in principle be measured in second order, within the wavelength range covered by the spectrometer. To improve on the relative accuracy of measurement of the two short-wavelength transitions, higher-order observations would be desirable, although in some higher orders, line blending will be a problem. (For example, the second order of the Mo XL $2s\ 2S_{1/2} - 2p\ 2P_{3/2}$ transition will fall in the wing of the strong resonance line of Mg-like Mo XXXI at $116.0\ \text{\AA}$)

III. RESULTS

The wavelengths obtained are $49.904 \pm 0.03\ \text{\AA}$ and $137.787 \pm 0.03\ \text{\AA}$ for the Mo XXXIX resonance and intercombination lines, respectively, while for Mo XL, the results are

58.499±0.02 Å and 143.998±0.02 Å for the $2s\ 2S_{1/2} - 2p\ 2P_{3/2}$ and $2s\ 2S_{1/2} - 2p\ 2P_{1/2}$ transitions, respectively. The corresponding energy level values are presented in Table I. Also given are the calculated energies by Johnson *et al.*¹ for Li-like Mo and the semi-empirical predictions by Edlén³ for Be-like Mo. Regarding Li-like Mo XL we shall here merely state that the observed deviation between experiment and theory is consistent with an observation in Kr⁷ and further down in the sequence. This deviation is attributed to a screening of the Lamb shift¹. For a detailed discussion of the lithium sequence we would like to refer to a recent paper by Hinnov, Denne *et al.*⁶. For the beryllium sequence resonance and intercombination lines, the present observations in Mo XXXIX confirm the systematic deviation from the semi-empirical predictions by Edlén³ as observed further down in the sequence by Denne and Hinnov². The difference between observed wavenumber and predicted by Edlén is illustrated in Fig. 3 for the resonance and intercombination lines for the range $Z=22-42$. The experimental wavelengths are given in Tables II and III for the $2s^2\ 1S_0 - 2s2p\ 1P_1$ and $2s^2\ 1S_0 - 2s2p\ 3P_1$ transitions, respectively. In order to obtain the predicted wavenumbers for elements beyond krypton, the formulae given in Ref. 3, Tables V and VI, were extrapolated. It is seen from the Figure that the resonance line wavelength is becoming increasingly longer than predicted (lower wavenumber) while the intercombination line wavelength is becoming increasingly shorter. In this context, it is interesting to note that the $1S_0 - 3P_1$ transition wavelength eventually becomes shorter than the $2S_{1/2} - 2P_{1/2}$ resonance line of the lithiumlike ion - this takes place near $Z=31$. In Cu ($Z=29$) the Cu XXVI $1S_0 - 3P_1$ line and the Cu XXVII $2S_{1/2} - 2P_{1/2}$ line are at 227.808 Å and 224.795 Å⁸, respectively, while in Ge ($Z=32$) the said transitions are at 199.36 Å⁹ and 200.290 Å⁶, respectively.

The deviation from Edlén's predictions is attributed to the lack of accurate data beyond Ni ($Z=28$) at the time (1983), and furthermore poor or conflicting measurements, particularly for iron and nickel. Since then, not only have measurements been extended to higher Z , but more precise measurements have become available for Fe and Ni. The most reliable and up-to-date experimental data to our knowledge are listed in Tables II and III, and have

been used in the comparison with Edlén's values. A weighed, least-squares fit of a polynomial in Z has been made to the difference between observed and predicted wavenumber, for both the resonance and the intercombination line. The fitted graphs are shown in Fig. 3. From the graphs we may obtain recalculated wavelengths to be compared with the measurements, as well as interpolated wavelengths for elements not yet investigated. Wavelengths thus obtained are given in the right-most column of Tables II and III. The accuracy of the recalculated wavelengths is estimated to be $\pm 0.02 \text{ \AA}$, or better.

The semi-empirical predictions by Edlén³ were based on multi-configuration Dirac-Fock calculations by Cheng, Kim and Desclaux¹¹. If we compare the observed wavenumbers with those of Cheng *et al.* in the Z -range shown in Fig. 3, we find that for the singlet line, the theoretical wavenumbers are too low by about 0.3 % throughout (*i.e.* the calculated wavelengths are too long). For the intercombination line, the theoretical wavenumbers are too low by about 0.2 % at Ni, and become too high for higher Z (about 0.15 % at $Z=42$) compared to the observed values. In view of the experimental accuracy now attainable, more accurate theoretical calculations are called for for the beryllium sequence, preferably on a par with those by Johnson *et al.* for the lithium isoelectronic sequence, although we appreciate the greater theoretical complexity associated with the presence of one more electron in the case of the berylliumlike ion.

ACKNOWLEDGEMENTS

The authors wish to thank Dr. E. Hinnov for several fruitful discussions.

The enthusiastic support of this experiment by the JET Operating Team, in particular Drs. M. Bures, P. Lallia and P. Nielsen, is gratefully acknowledged.

REFERENCES

1. W. R. Johnson, S. A. Blundell, and J. Sapirstein, *Phys. Rev. A* **37**, 2764 (1988).
2. B. Denne and E. Hinnov, *Phys. Scripta* **35**, 811 (1987).
3. B. Edlén, *Phys. Scripta* **28**, 51 (1983).
4. G. Magyar, M. R. Barnes, S. Cohen, A. Edwards, N. C. Hawkes, D. Pasini, and P. J. Roberts, JET Report JET-R(88)15 (1988).
5. J.-L. Schwob, A. W. Wouters, S. Suckewer, and M. Finkenthal, *Rev. Sci. Instrum.* **58**, 1601 (1987).
6. E. Hinnov and the TFTR Operating Team, and B. Denne and the JET Operating Team (to be published).
7. B. Denne, E. Hinnov, J. Ramette, and B. Saoutic (to be published).
8. E. Hinnov (private communication).
9. E. Hinnov, F. Boody, S. Cohen, U. Feldman, J. Hosea, K. Sato, J.-L. Schwob, S. Suckewer, and A. Wouters, *J. Opt. Soc. Am.* **B3**, 1288 (1986).
10. E. Hinnov, A. Ramsey, B. Stratton, S. Cohen, and J. Timberlake, *J. Opt. Soc. Am.* **B4**, 1293 (1987).
11. K. T. Cheng, Y.-K. Kim, and J. P. Desclaux, *At. Data and Nucl. Data Tables* **24**, 111 (1979).

TABLE I. Energy levels (in cm^{-1}) of Mo XXXIX and Mo XL.

Spectrum	Level	Experimental energy	Predicted energy
Mo XXXIX	$2s^2\ ^1S_0$	0	0
	$2s2p\ ^3P_1$	725758 ± 158	722026^a
	$\ ^1P_1$	2003847 ± 1200	2009300^a
Mo XL	$2s\ ^2S_{1/2}$	0	0
	$2p\ ^2P_{1/2}$	694454 ± 96	715788^b
	$\ ^2P_{3/2}$	1709430 ± 585	1729001^b

^aRef. 3 (extrapolated).

^bRef. 1 (interpolated).

TABLE II. Observed wavelength (λ_o) and wavenumber (σ_o) for the $2s^2\ ^1S_0 - 2s2p\ ^1P_1$ transition of the Be-sequence for elements in the range $Z=22-42$. σ_E is the wavenumber predicted by Edlén, and λ_c is the wavelength obtained from the graph in Fig. 3.

Z	Spectrum	λ_o (Å)	σ_o (cm ⁻¹)	σ_E (cm ⁻¹)	λ_c (Å)
22	Ti XIX	169.585±0.02 ^a	589675± 70	589710	169.572
23	V XX			627555	159.362
24	Cr XXI	149.90 ±0.03 ^b	667111± 135	667166	149.913
25	Mn XXII			708759	141.126
26	Fe XXIII	132.913±0.01 ^b	752372± 57	752574	132.920
27	Co XXIV			798863	125.227
28	Ni XXV	117.995±0.01 ^a	847494± 72	847894	117.996
29	Cu XXVI	111.186±0.01 ^b	899394± 81	899954	111.180
30	Zn XXVII			955349	104.744
31	Ga XXVIII			1014397	98.659
32	Ge XXIX	92.90 ±0.03 ^a	1076426± 350	1077432	92.898
33	As XXX			1144793	87.444
34	Se XXXI	82.285±0.010 ^b	1215288± 150	1216835	82.280
35	Br XXXII			1293933	77.390
36	Kr XXXIII	72.756±0.02 ^c	1374457± 380	1376480	72.761
37	Rb XXXIV			1464876	68.383
38	Sr XXXV			1559524	64.245
39	Y XXXVI			1660847	60.337
40	Zr XXXVII			1769281	56.651
41	Nb XXXVIII			1885280	53.176
42	Mo XXXIX	49.904±0.03 ^d	2003847±1200	2009300	49.904

^aRef. 2.

^bRef. 8.

^cRef. 7.

^dPresent work.

TABLE III. As Table II for the $2s^2\ ^1S_0 - 2s2p\ ^3P_1$ transition for elements in the range $Z=26-42$.

Z	Spectrum	λ_o (Å)	σ_o (cm ⁻¹)	σ_E (cm ⁻¹)	λ_c (Å)
26	Fe XXIII	263.755±0.01 ^a	379140± 20	379164	263.738
27	Co XXIV			398737	250.777
28	Ni XXV	238.83 ±0.02 ^b	418708± 35	418653	238.836
29	Cu XXVI	227.808±0.01 ^a	438966± 19	438896	227.808
30	Zn XXVII			459443	217.604
31	Ga XXVIII			480273	208.144
32	Ge XXIX	199.360±0.03 ^c	501605± 75	501362	199.360
33	As XXX			522686	191.190
34	Se XXXI	183.580±0.02 ^d	544722± 60	544223	183.580
35	Br XXXII			565948	176.480
36	Kr XXXIII	169.845±0.025 ^e	588772± 87	587844	169.846
37	Rb XXXIV			609899	163.635
38	Sr XXXV			632091	157.812
39	Y XXXVI			654410	152.345
40	Zr XXXVII			676845	147.202
41	Nb XXXVIII			699390	142.357
42	Mo XXXIX	137.787±0.03 ^f	725758±158	722026	137.787

^aRef. 8.

^bRef. 2.

^cRef. 9.

^dRef. 10.

^eRef. 7.

^fPresent work.




Fig. 1 Spectrum in the region $\sim 49-59 \text{ \AA}$ before (*dashed line*) and after (*solid line*) injection of molybdenum.




Fig. 2 Time evolution of the Mo_{xxxii} 127.868 \AA line (Na-like) and the Li-like Mo_{xl} line at 143.998 \AA . Also shown (*top*) is the central electron temperature, $T_e(0)$.




Fig. 3 Difference between observed wavenumber (σ_0) and predicted by Edlén (σ_E) for the Be-sequence resonance line (filled circles) and intercombination line (open circles) in the range $Z = 22-42$. The graphs are least-squares polynomial fits to the respective data points.

APPENDIX 1.

THE JET TEAM

JET Joint Undertaking, Abingdon, Oxon, OX14 3EA, U.K.

J. M. Adams¹, F. Alladio⁴, H. Altmann, R. J. Anderson, G. Appruzzese, W. Bailey, B. Balet, D. V. Bartlett, L. R. Baylor²⁴, K. Behringer, A. C. Bell, P. Bertoldi, E. Bertolini, V. Bhatnagar, R. J. Bickerton, A. Boileau³, T. Bonicelli, S. J. Booth, G. Bosia, M. Botman, D. Boyd³¹, H. Brelen, H. Brinkschulte, M. Brusati, T. Budd, M. Bures, T. Businaro⁴, H. Buttgereit, D. Cacaut, C. Caldwell-Nichols, D. J. Campbell, P. Card, J. Carwardine, G. Celentano, P. Chabert²⁷, C. D. Challis, A. Cheetham, J. Christiansen, C. Christodoulopoulos, P. Chuilon, R. Claesen, S. Clement³⁰, J. P. Coad, P. Colestock⁶, S. Conroy¹³, M. Cooke, S. Cooper, J. G. Cordey, W. Core, S. Corti, A. E. Costley, G. Cottrell, M. Cox⁷, P. Cripwell¹³, F. Crisanti⁴, D. Cross, H. de Blank¹⁶, J. de Haas¹⁶, L. de Kock, E. Deksnis, G. B. Denne, G. Deschamps, G. Devillars, K. J. Dietz, J. Dobbing, S. E. Dorling, P. G. Doyle, D. F. Düchs, H. Duquenoy, A. Edwards, J. Ehrenberg¹⁴, T. Elevant¹², W. Engelhardt, S. K. Erents⁷, L. G. Eriksson⁵, M. Evrard², H. Falter, D. Flory, M. Forrest⁷, C. Froger, K. Fullard, M. Gadeberg¹¹, A. Galetsas, R. Galvao⁸, A. Gibson, R. D. Gill, A. Gondhalekar, C. Gordon, G. Gorini, C. Gormezano, N. A. Gottardi, C. Gowers, B. J. Green, F. S. Grigh, M. Gryzinski²⁶, R. Haange, G. Hammett⁶, W. Han⁹, C. J. Hancock, P. J. Harbour, N. C. Hawkes⁷, P. Haynes⁷, T. Hellsten, J. L. Hemmerich, R. Hemsworth, R. F. Herzog, K. Hirsch¹⁴, J. Hoekzema, W. A. Houlberg²⁴, J. How, M. Huart, A. Hubbard, T. P. Hughes³², M. Hugon, M. Huguet, J. Jacquinet, O. N. Jarvis, T. C. Jernigan²⁴, E. Joffrin, E. M. Jones, L. P. D. F. Jones, T. T. C. Jones, J. Källne, A. Kaye, B. E. Keen, M. Keilhacker, G. J. Kelly, A. Khare¹⁵, S. Knowlton, A. Konstantellos, M. Kovanen²¹, P. Kupschus, P. Lallia, J. R. Last, L. Lauro-Taroni, M. Laux³³, K. Lawson⁷, E. Lazzaro, M. Lennholm, X. Litaudon, P. Lomas, M. Lorentz-Gottardi², C. Lowry, G. Magyar, D. Maisonnier, M. Malacarne, V. Marchese, P. Massmann, L. McCarthy²⁸, G. McCracken⁷, P. Mendonca, P. Meriguet, P. Micozzi⁴, S. F. Mills, P. Millward, S. L. Milora²⁴, A. Moissonnier, P. L. Mondino, D. Moreau¹⁷, P. Morgan, H. Morsi¹⁴, G. Murphy, M. F. Nave, M. Newman, L. Nickesson, P. Nielsen, P. Noll, W. Obert, D. O'Brien, J. O'Rourke, M. G. Pacco-Düchs, M. Pain, S. Papastergiou, D. Pasini²⁰, M. Paume²⁷, N. Peacock⁷, D. Pearson¹³, F. Pegoraro, M. Pick, S. Pitcher⁷, J. Plancoulaine, J-P. Poffé, F. Porcelli, R. Prentice, T. Raimondi, J. Ramette¹⁷, J. M. Rax²⁷, C. Raymond, P-H. Rebut, J. Removille, F. Rimini, D. Robinson⁷, A. Rolfe, R. T. Ross, L. Rossi, G. Rupprecht¹⁴, R. Rushton, P. Rutter, H. C. Sack, G. Sadler, N. Salmon¹³, H. Salzmann¹⁴, A. Santagiustina, D. Schissel²⁵, P. H. Schild, M. Schmid, G. Schmidt⁶, R. L. Shaw, A. Sibley, R. Simonini, J. Sips¹⁶, P. Smeulders, J. Snipes, S. Sommers, L. Sonnerup, K. Sonnenberg, M. Stamp, P. Stangeby¹⁹, D. Start, C. A. Steed, D. Stork, P. E. Stott, T. E. Stringer, D. Stubberfield, T. Sugie¹⁸, D. Summers, H. Summers²⁰, J. Taboda-Duarte²², J. Tagle³⁰, H. Tamnen, A. Tanga, A. Taroni, C. Tebaldi²³, A. Tesini, P. R. Thomas, E. Thompson, K. Thomsen¹¹, P. Trevalion, M. Tschudin, B. Tubbing, K. Uchino²⁹, E. Usselmann, H. van der Beken, M. von Hellermann, T. Wade, C. Walker, B. A. Wallander, M. Walravens, K. Walter, D. Ward, M. L. Watkins, J. Wesson, D. H. Wheeler, J. Wilks, U. Willen¹², D. Wilson, T. Winkel, C. Woodward, M. Wykes, I. D. Young, L. Zannelli, M. Zarnstorff⁶, D. Zsche¹⁴, J. W. Zwart.

PERMANENT ADDRESS

1. UKAEA, Harwell, Oxon. UK.
2. EUR-EB Association, LPP-ERM/KMS, B-1040 Brussels, Belgium.
3. Institute National des Recherches Scientifique, Quebec, Canada.
4. ENEA-CENTRO Di Frascati, I-00044 Frascati, Roma, Italy.
5. Chalmers University of Technology, Göteborg, Sweden.
6. Princeton Plasma Physics Laboratory, New Jersey, USA.
7. UKAEA Culham Laboratory, Abingdon, Oxon. UK.
8. Plasma Physics Laboratory, Space Research Institute, Sao José dos Campos, Brazil.
9. Institute of Mathematics, University of Oxford, UK.
10. CRPP/EPFL, 21 Avenue des Bains, CH-1007 Lausanne, Switzerland.
11. Risø National Laboratory, DK-4000 Roskilde, Denmark.
12. Swedish Energy Research Commission, S-10072 Stockholm, Sweden.
13. Imperial College of Science and Technology, University of London, UK.
14. Max Planck Institut für Plasmaphysik, D-8046 Garching bei München, FRG.
15. Institute for Plasma Research, Gandhinagar Bhat Gujrat, India.
16. FOM Instituut voor Plasmafysica, 3430 Be Nieuwegein, The Netherlands.
17. Commissariat à l'Energie Atomique, F-92260 Fontenay-aux-Roses, France.
18. JAERI, Tokai Research Establishment, Tokai-Mura, Naka-Gun, Japan.
19. Institute for Aerospace Studies, University of Toronto, Downsview, Ontario, Canada.
20. University of Strathclyde, Glasgow, G4 ONG, U.K.
21. Nuclear Engineering Laboratory, Lapeenranta University, Finland.
22. JNICT, Lisboa, Portugal.
23. Department of Mathematics, Univeristy of Bologna, Italy.
24. Oak Ridge National Laboratory, Oak Ridge, Tenn., USA.
25. G.A. Technologies, San Diego, California, USA.
26. Institute for Nuclear Studies, Swierk, Poland.
27. Commissariat à l'Energie Atomique, Cadarache, France.
28. School of Physical Sciences, Flinders University of South Australia, South Australia 5042.
29. Kyushi University, Kasagu Fukuoka, Japan.
30. Centro de Investigaciones Energeticas Medioambientales y Techalogicas, Spain.
31. University of Maryland, College Park, Maryland, USA.
32. University of Essex, Colchester, UK.
33. Akademie de Wissenschaften, Berlin, DDR.