



EUROfusion

WP17ER-PR(18) 20014

A Kleiner et al.

**Free boundary 3D ideal MHD
equilibrium calculations for non-linearly
saturated current driven external kink
modes in tokamaks**

Preprint of Paper to be submitted for publication in
Nuclear Fusion Letter



This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

This document is intended for publication in the open literature. It is made available on the clear understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EUROfusion Programme Management Unit, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK or e-mail Publications.Officer@euro-fusion.org

Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EUROfusion Programme Management Unit, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK or e-mail Publications.Officer@euro-fusion.org

The contents of this preprint and all other EUROfusion Preprints, Reports and Conference Papers are available to view online free at <http://www.euro-fusionscipub.org>. This site has full search facilities and e-mail alert options. In the JET specific papers the diagrams contained within the PDFs on this site are hyperlinked

Free boundary 3D ideal MHD equilibrium calculations for non-linearly saturated current driven external kink modes in tokamaks

A Kleiner¹, J P Graves¹, W A Cooper¹, T Nicolas² and C Wahlberg³

¹ Ecole Polytechnique Fédérale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015 Lausanne, Switzerland

² Centre de Physique Théorique, Ecole Polytechnique, CNRS, F-91128 Palaiseau Cedex, France

³ Department of Physics and Astronomy, PO Box 516, Uppsala University, SE-751 20 Uppsala, Sweden

E-mail: andreas.kleiner@epfl.ch

Abstract. It is shown that free boundary 3D equilibrium calculations in tokamak geometry are capable of capturing the physics of non-linearly saturated external kink modes for monotonic current and q profiles typical of standard (baseline) plasma scenarios. The VMEC ideal MHD equilibrium model exhibits strong flux surface corrugations of the plasma vacuum boundary, driven by the core current profile. A method is presented which conveniently extracts the amplitude of the corrugation in terms of Fourier components in straight field line coordinates. The Fourier spectrum, and condition for non-linear corrugation agrees well with linear simulations, and the saturated amplitude agrees well with non-linear analytic calculations.

PACS numbers: 52.25.Xz, 52.30.Cv, 52.55.-s, 52.55.Fa

Keywords: MHD, external kink, tokamak, free boundary, 3D equilibrium, nonlinear

Submitted to: *Nucl. Fusion*

1. Introduction

External kink modes are known [1, 2] to be of concern for the development of plasma scenarios. They set operational limits and define inaccessible windows for the edge safety factor and the peaking of the current profile. External kink modes tend to be linearly unstable for large current gradients, especially when these gradients are close to the plasma edge. After an initial linear growth in the non-linear phase, a mode usually saturates (non-linear stability) [3, 4], though in practice, the corrugation can be so large as to touch the vacuum vessel or plasma-facing components, thus causing disruptions [1].

The investigation of non-linear instabilities is typically carried out [5] with initial value codes (e.g. XTOR-2F [6]). Such simulations are computationally expensive due to the resolution and number of time steps necessary and require a certain level of parallelisation [7]. Since a saturated state without equilibrium flows is characterised by being time-independent, an equilibrium code can in theory obtain a saturated state, satisfying the force balance equation $\mathbf{j} \times \mathbf{B} - \nabla p = 0$. Indeed, this was already shown for non-resonant internal kink modes with $q > 1$, where initial value (XTOR-2F) saturated states and VMEC fixed-boundary calculations showed similar $m = n = 1$ displacement amplitudes [5].

The goal of this paper is to investigate the degree to which 3D free-boundary equilibrium codes can model conventional external kink modes, driven by current gradients associated with standard tokamak operation (baseline scenario) with monotonic (standard [Wesson-like [2]]) q profiles. The work differs from recent [8, 9, 10] deployment of VMEC for modelling edge harmonic oscillations in plasmas with low magnetic shear at the edge. Those oscillations are driven predominantly by pressure gradients, rather than current gradients. The 3D converged equilibria in this manuscript will be compared with the spectral properties of linear numerical stability analysis, and further verified by analytic non-linear calculations [4]. Lack of a converged equilibrium will be taken to mean that an unstable external kink mode forms, but non-linear saturation has not been achieved. The advantages of the equilibrium approach to external kink mode problems are numerous. First, the computations are fast and accurate, and second they provide convenient Fourier decomposed magnetic geometry for advanced studies such as fast particle [11] and impurity transport [12] in non-axisymmetric plasmas. Such studies require long and highly accurate particle simulations. To obtain the required accuracy, the VENUS-LEVIS guiding center code [13] for example exploits the Fourier decomposition of the magnetic field for orbit calculations.

In this letter we present ideal MHD free-boundary computations of JET-like plasmas carried out with

the VMEC code [14]. First, the characteristics of the observed three-dimensional equilibria and the choice of the coordinate system for mode spectrum analysis are discussed. This is followed by the VMEC calculation of the non-linear saturated displacement amplitude for varying edge q value, and two choices of current density profiles. The results are then compared with the helical external kink amplitude calculated from an analytical model based on a non-linear large aspect ratio expansion.

2. Non-axisymmetric VMEC equilibria

The computation of 3D ideal MHD equilibria is performed using the free-boundary VMEC code [14, 15], which arrives at an equilibrium state by minimising the energy functional

$$W = \iiint_{\text{plasma}} \left[\frac{|\mathbf{B}|^2}{2\mu_0} + \frac{p(\rho)}{\Gamma - 1} \right] d^3x - \iiint_{\text{vacuum}} \frac{|\mathbf{B}_v|^2}{2\mu_0} d^3x, \quad (1)$$

where \mathbf{B} is the magnetic field inside the plasma, \mathbf{B}_v is the vacuum magnetic field, $p(\rho)$ is the pressure as a function of the radial variable $\rho = \sqrt{\Phi/\Phi_a}$. The toroidal flux is denoted by Φ , and subscript ' a ' indicates that a quantity is evaluated at the plasma boundary. The poloidal and toroidal magnetic field coils are modeled as discretised filaments with specified coil currents. In this proof-of-principle study we employ an accurate model of the up-down symmetric toroidal and poloidal JET coils [10], but we do not include the effects of other coil systems (e.g. divertor coils), the vacuum vessel conductor nor iron core. The vacuum magnetic field is calculated from the coil currents according to the Biot-Savart law [16]. During the iterations that minimise the energy functional with respect to an artificial time variable, the plasma boundary is free to move. We impose stellarator symmetry on the 3D equilibrium and choose 289 flux surfaces with a mode spectrum of $0 \leq m \leq 14$ poloidal and $-6 \leq n \leq 6$ toroidal modes. In order to converge towards the 3D equilibrium quickly, we provide a small perturbation to the magnetic axis initial guess.

The linear growth rate and the non-linear saturated amplitude of external kink modes depend on the distance to a conducting wall surrounding the plasma. The VMEC model does not include such a wall. Instead, the plasma is enclosed inside a structure of magnetic field coils at a finite distance from the plasma. We hence cannot investigate the influence of the wall distance on the VMEC edge corrugations. For comparison with VMEC, the conducting wall distance must be set to infinity (very large) for models that have the effect of a conducting wall. These models (linear and non-linear) will investigate the sensitivity of wall distance, and thus the likely correction to VMEC due to the vacuum vessel.




Figure 1: Plasma boundary and magnetic axis (indicated by a cross) of a free-boundary VMEC equilibrium with $q_a = 3.752$ and $\beta_N = 0.552\%$ are shown at different toroidal angles ϕ . The non-axisymmetric edge corrugation is clearly visible.

We start by examining dominant $n = 1, m = 4$ external modes with edge safety factor $q_a \lesssim 4$. Following the philosophy of previous studies of non-linear external kink modes [4] we choose a current density profile $j(\rho) \propto (1 - \rho^8)$, which is similar to those used by Wesson [2], however with a steeper gradient to make the non-linear structure in VMEC simulations larger, and hence easier to resolve numerically. For all simulations the pressure is chosen to be $p(\rho) \propto (1 - \rho^2)$. In order to investigate the effect of finite pressure on the resulting edge corrugations, $p(\rho)$ is multiplied by a scalar to achieve different values of $\beta_N = \beta a B_\phi / I_p$, where a is the minor radius, B_ϕ the toroidal magnetic field and I_p the plasma current. The value of q_a (edge safety factor) is crucial for both linear and non-linear external kink stability. We vary I_p in the VMEC simulations in order to study the dependency of the saturated edge displacement η on q_a with the given forms of $j(\rho)$ and varying pressure. The q profiles resulting from these equilibrium configurations are monotonic where $q(\rho) \approx \frac{B_\phi \rho^2}{\mu_0 R} / \int_0^\rho j(\rho') \rho' d\rho'$.

First, we simulate a plasma with a total toroidal current of $I_t = 2.6$ MA such that the resulting edge safety factor of $q_a = 3.752$ is below the rational value $m/n = 4/1$. The resulting VMEC equilibrium is non-axisymmetric with a strong edge corrugation in the toroidal and poloidal directions. This is illustrated in Fig. 1, where the plasma boundary and the magnetic axis are shown at different toroidal angles ϕ . From Fig. 2a, which shows a poloidal cross section of the flux surfaces for the $\phi = 0$ case of Fig. 1, it is evident that the perturbation is strongest at the plasma boundary and decreases towards the magnetic axis where it vanishes (hence only one cross observed in Fig. 1). This behaviour as expected is characteristic of external




Figure 2: Visualisation of the resulting 3D equilibrium with $q_a = 3.752$ shown for the $\phi = 0$ case of Fig. 1. (a) Shape and position of the magnetic flux surfaces. The perturbation is strongest at the plasma edge and vanishes towards the magnetic axis. (b) Perturbation $\delta p = p_{3D} - p_{2D}$ arising from the comparison of the pressure in the 3D equilibrium with the neighbouring state.

kink modes. By retaining only $n = 0$ modes, we can find a neighbouring axisymmetric state to the obtained 3D equilibrium. A comparison of 3D VMEC states with their neighbouring axisymmetric equilibria was previously used to determine toroidal field ripple [17]. Defining $\delta p = p_{3D} - p_{2D}$ as being the difference of p in the 3D state compared to the axisymmetric state, we can identify the periodicity and the location of the perturbation. It is noted in passing that the 3D displacement ξ of the magnetic surfaces can be approximately determined via $\xi = -\delta p (\partial p_{2D} / \partial \rho)^{-1}$. Fig. 2b shows δp for the $\phi = 0$ case of Fig. 1. It can be seen that the perturbation is located at the plasma edge and the $m = 4$ character is evident.

To determine the non-linear amplitude, emphasis has to be placed on the coordinate system underlying the spectral analysis of the displacement. From VMEC, the axisymmetric flux surface $\mathbf{S}_{2D}(\theta, \phi)$ as well as 3D flux surface $\mathbf{S}_{3D}(\theta, \phi)$ are known in VMEC flux coordinates $(s = \rho^2, \theta, \phi)$ [14]. For a given point \mathbf{P}_0 lying on \mathbf{S}_{2D} with normal vector \mathbf{N} , a convenient definition of the corrugation displacement at the edge is

$$\eta(\theta, \phi) \mathbf{N}(\theta, \phi) = \mathbf{S}_{3D} - \mathbf{P}_0(\theta, \phi). \quad (2)$$

The Fourier series representing η is of the same form as that of the magnetic field strength, i.e.,

$$\eta(\theta, \phi) = \sum_{m=0}^{M_{\max}} \sum_{n=0}^{N_{\max}} \eta_{mn} \cos(m\theta - n\phi), \quad (3)$$

due to the imposed stellarator symmetry. We can calculate the mode spectrum separately for $\eta(\theta, \phi)$,




Figure 3: Mode spectrum of the edge displacement obtained in two different coordinate systems. (a) In VMEC coordinates. (b) In straight field line coordinates (dominant $m = 4/n = 1$ mode).

$|\mathbf{B}|(\theta, \phi)$ as well as $R(\theta, \phi)$ and $Z(\theta, \phi)$ at the edge. The spectrum of each of these quantities is rich and exhibits a variety of strong poloidal modes for $n = 1$. For the case shown in Fig. 2, the 4/1 VMEC coordinate coefficients of R , Z , $|\mathbf{B}|$ and η are not dominant. This is in contradiction with the visual observations from the magnetic topology, which has a clear 4/1 structure. To allow a direct comparison with η obtained in straight field line (sfl) coordinates, the mode amplitudes for $\eta(\phi, \theta)$ in VMEC coordinates can be seen in Fig. 3a.

Mode spectra are expected to be most narrow for a straight field line coordinate system. Dominant poloidal modes can thus be identified for comparison with those seen in Figs. 1 and 2 (simply by inspection), and for suitable comparison with the non-linear analytic treatment described later. For this we need to transform the poloidal angle of the VMEC coordinate system to a straight field line angle θ_{sfl} by equating the volume elements in VMEC coordinates and straight field line coordinates

$$\mathcal{J}_{\text{sfl}} d\phi d\theta_{\text{sfl}} d\rho^2 = \mathcal{J}_{\text{VMEC}} d\phi d\theta d\rho^2. \quad (4)$$

Since the toroidal angle ϕ and the radial variable ρ are identical in both coordinate systems, the only variables remaining are θ_{sfl} and θ . Integration of Eq. (4) yields

$$\theta_{\text{sfl}}(\theta) = 2\pi \left[\int_0^{2\pi} \frac{\mathcal{J}_{\text{VMEC}}(\theta)}{R^2(\theta)} d\theta \right]^{-1} \int_0^\theta \frac{\mathcal{J}_{\text{VMEC}}(\theta)}{R^2(\theta)} d\theta, \quad (5)$$

where $\mathcal{J}_{\text{sfl}} \propto R^2$ was used and $\mathcal{J}_{\text{VMEC}}$ and R are computed by VMEC from the axisymmetric equilibrium. Now with Eq. (2) we are able to calculate




Figure 4: (a) Linear external kink growth rate γ of the 4/1 mode calculated with KINX. (b) Linear $n = 1$ (KINX) radial displacement functions for various poloidal mode numbers (sfl coordinates). (c) Non-linear $n = 1$ saturated (VMEC) radial displacement functions η for various poloidal mode numbers (sfl coordinates).

the coefficients for the Fourier series of $\eta(\theta_{\text{sfl}}, \phi)$. In this sfl coordinate system the mode spectrum cleanly and clearly identifies a standard external kink mode. As shown in Fig. 3b the 4/1 mode is dominant as expected from the visual observations of Fig. 2 as well as from external kink mode theory. With all other modes being fairly weak compared to the dominant one, poloidal and toroidal coupling thus appears to have a weak effect. Toroidicity and beta can be expected to have only a mild effect on mode structure, and hence we can compare the mode amplitudes to an analytical model that assumes cylindrical geometry [4].

To obtain non-linearly saturated external kink modes, the system is initially required to be linearly unstable. This condition is verified from numerical computations with the linear eigenvalue code KINX [18]. The effects of small aspect ratio, shaping and finite beta are retained in these computations and in Figs. 4a, 4b the distance to the perfectly conducting wall surrounding the plasma is set to $b/a = 10$, i.e.

sufficiently large to consider no effects due to the wall. The linear growth rate $\gamma = -i\omega$ of the 4/1 mode is calculated as a function of q_a . This is presented in Fig. 4a, where γ is normalised by the Alfvén frequency ω_A . The 4/1 external kink mode is linearly unstable in a wide range of q_a , where saturated non-linear states can possibly arise. The linear radial eigenfunctions of the sfl harmonics in KINX resemble the mode spectrum calculated in sfl coordinates in Fig. 3b (for the non-linear equilibrium calculations). For an edge safety factor below $q_a < 4$, the eigenfunction (radial displacement) of the 4/1 mode is dominant at the plasma boundary. This is shown in Fig. 4b for the case with an edge safety factor of $q_a = 3.752$. The similarity of these linear eigenfunctions and the non-linear saturated VMEC displacement η_{mn} as a function of the radial variable, shown in Fig. 4c, is remarkable.

3. Comparison with saturated external kink mode amplitude

After demonstrating the external kink spectral properties of the VMEC 3D equilibria, we now compare the non-linear saturated amplitude of the displacement in VMEC with analytical predictions. In the following we provide a concise and self-contained summary of an analytical model [4] for the non-linear saturated external kink amplitude. In this context, we correct Eq. (34) of [4], where one term was found to be missing. The model assumes cylindrical geometry, circular cross section and no pressure effects. For non-resonant modes the non-linear evolution [19, 20] of the helical displacement η is given by the solution of

$$\frac{\partial^2 \eta}{\partial t^2} + D_1 \eta + D_3 \eta^3 = 0, \quad (6)$$

where the parameters D_1 and D_3 are defined below. For the saturated and hence time-independent state the solution to the amplitude η is

$$\eta = \sqrt{-\frac{D_1}{D_3}}. \quad (7)$$

For linear instability D_1 is negative. It is proportional to the square of the linear growth rate $-\omega^2$ and thus $D_1 = 0$ defines the marginal points. Non-linear stability is determined by the coefficient D_3 . The system is non-linearly stable, i.e. the mode can saturate if $D_3 > 0$. We do not explicitly explore conditions of non-linear instability where $D_3 < 0$. The coefficients D_1 and D_3 are obtained from physical constraints on the plasma surface [4] and read

$$D_1 = [F(f_{1\rho} - m^2 \beta_1 + 1) - 2] F, \quad (8)$$

and

$$\begin{aligned} D_3 = & \frac{F^2 f_{1\rho}}{4} [6f_{1\rho}(f_{1\rho} + 1) + m^2 - 3] \\ & - F^2 \left(\frac{3}{2} n_{2\rho} f_{1\rho} - n_{3\rho} + \frac{1}{4} \right) \\ & + m^2 F^2 \left[\frac{\beta_1}{4} (6m^2 + 1) + m^2 \beta_1 \beta_2 (1 - 2m^2 \beta_1) \right. \\ & \left. - \frac{11}{8} m^2 \beta_1^2 + \frac{3}{8} \right] + 2n_2 + \frac{1}{\ln b} (Fm^2 \beta_1 + 1)^2 \quad (9) \\ & - F \left[m^2 \left(\frac{7}{2} \beta_1 - \beta_2 + 4m^2 \beta_1 \beta_2 - 2 \right) - \frac{1}{2} \right] \\ & + \frac{n_2 F^2}{2} [m^2 (\beta_1 (m^2 \beta_1 - 1) + 8m^2 \beta_1 \beta_2 - 5) + 1] \\ & + n_2 F [2m^2 (\beta_1 + 2\beta_2) - 1] - 2m^2 \beta_2 - \frac{5}{2}, \end{aligned}$$

where $x_\rho = \partial x / \partial \rho$. The coefficients D_1 and D_3 in Eqs. (8) and (9) differ from D_1 and D_3 in Eq. (6) by a constant but common factor that cancels out in the calculation of η in Eq. (7). The wall distance b is normalised by the minor radius, i.e. for $b = 1$ the wall is located at the plasma boundary and we define $\beta_1 := -\frac{1}{m} \frac{b^{2m} + 1}{b^{2m} - 1}$ and $\beta_2 := -\frac{1}{2m} \frac{b^{4m} + 1}{b^{4m} - 1}$. The function F is defined as

$$F(\rho) := \frac{I(\rho)}{\rho^2} - \frac{nq_a}{m}, \quad (10)$$

where $I(\rho) = \int_0^\rho 2\rho' j(\rho') d\rho'$ is the total current enclosed by a flux surface with radius ρ and normalised such that $I(1) = 1$. To obtain the functions f_1 , n_2 and n_3 , the helical amplitude η is expanded up to third order and a coupled system of differential equations arising from the ideal MHD model is solved numerically providing a solution for η to each order. These solutions can be written as sums of homogeneous $f_i(\rho)$ and particular $g_i(\rho)$ solutions and are denoted $\eta_m^{(1)}(\rho) \equiv f_1(\rho)\eta_m^{(1)}(1)$ to first, $\eta_{2m}^{(2)}(\rho) \equiv f_2(\rho)\eta_{2m}^{(2)}(1) + g_2(\rho)$ to second and $\eta_m^{(3)}(\rho) \equiv g_3(\rho)$ to third order. They satisfy the boundary conditions

$$\begin{aligned} \lim_{\rho \rightarrow 0} f_1(\rho) & \propto \rho^{m-1}, & f_1(1) & = 1, \\ \lim_{\rho \rightarrow 0} f_2(\rho) & \propto \rho^{2m-1}, & f_2(1) & = 1, \\ g_2(0) = g_2(1) & = 0, & g_3(0) = g_3(1) & = 0. \end{aligned} \quad (11)$$

Using a normalisation constant $\hat{\eta} := \eta_m^{(1)}(1)$, we further define the normalised solutions to second and third order $n_2 := \eta_{2m}^{(2)}(\rho) / \hat{\eta}^2$ and $n_3 := g_3(\rho) / \hat{\eta}^3$ respectively. In Eqs. (8) and (9), $F, f_{1\rho}, n_2, n_{2\rho}$ and $n_{3\rho}$ are evaluated at $\rho = 1$. Defining the operator

$$\mathcal{L}_k = \frac{d^2}{d\rho^2} + \left[\frac{3}{\rho} + 2 \frac{d}{d\rho} \ln F(\rho) \right] \frac{d}{d\rho} + \frac{1 - k^2 m^2}{\rho^2}, \quad (12)$$

with $k = 1, 2$ we obtain the solutions for η to each order from the following system of ODEs: [4]

$$\mathcal{L}_1 \eta_m^{(1)} = 0, \quad (13)$$

$$\mathcal{L}_2 \eta_{2m}^{(2)} = -\frac{F_\rho}{2F\rho^2} \left(3\rho^2 (\eta_{m\rho}^{(1)})^2 + m^2 (\eta_m^{(1)})^2 \right) - \frac{1}{2\rho^3} \left[(3m^2 - 1) (\eta_m^{(1)})^2 + 2\rho \eta_m^{(1)} \eta_{m\rho}^{(1)} + 5\rho^2 (\eta_{m\rho}^{(1)})^2 \right], \quad (14)$$

$$\begin{aligned} \mathcal{L}_1 \eta_m^{(3)} &= (\eta_m^{(1)})^3 \frac{1}{4\rho^4 F} [F(12m^2 - m^4 - 7) - 2\rho m^2 F_\rho] \\ &+ \eta_{m\rho}^{(1)} (\eta_m^{(1)})^2 \frac{1}{4\rho^3 F} [F(2 - 16m^2) - 3\rho m^2 F_\rho] + (\eta_{m\rho}^{(1)})^2 \eta_m^{(1)} \frac{1}{4\rho^2 F} [F(41 - 19m^2) + 12\rho F_\rho] \\ &+ \eta_m^{(1)} \eta_{2m}^{(2)} \frac{1}{\rho^3 F} [F(1 - 3m^2) + 2\rho m^2 F_\rho] - \eta_m^{(1)} \eta_{2m\rho}^{(2)} \frac{1}{\rho^2} (1 - 3m^2) \\ &+ (\eta_{m\rho}^{(1)})^3 \frac{1}{4\rho F} [18F + 15\rho F_\rho] - \eta_{m\rho}^{(1)} \eta_{2m\rho}^{(2)} \frac{1}{\rho F} [5F + 3\rho F_\rho] - \eta_{m\rho}^{(1)} \eta_{2m}^{(2)} \frac{1}{\rho^2} (1 - 6m^2). \end{aligned} \quad (15)$$

The normalisation constant $\hat{\eta}$ can be eliminated by exploiting pressure balance across the plasma surface [4], resulting in the expression

$$n_2(1) = \frac{-\frac{1}{4} \left(m^2 (m^2 \beta_1^2 + 8m^2 \beta_1 \beta_2 - 4\beta_2 - 2) - 3(f_{1\rho})^2 + 4 \frac{g_{2\rho}^2}{\hat{\eta}^2} + 1 \right) F^2 - \left(m^2 (\beta_1 + 2\beta_2) - \frac{1}{2} \right) F - 1}{(F(f_{2\rho} - 4m^2 \beta_2 + 1) - 2) F}. \quad (16)$$




Figure 5: Dependence of the saturated 4/1 external kink amplitude on the wall distance b as calculated from the non-linear analytical model.

We stress the fact that in this model, η depends only on single helical mode numbers m and n (as expected for cylindrical geometry), current density profile $j(\rho)$, wall distance b and edge safety factor q_a . In order to study the degree of relevance of the equilibrium free-boundary approach without a conducting wall, it is important to investigate the sensitivity of the effect of the conducting wall distance from the plasma on the non-linear saturated state. This is achieved with the analytical model. Figure 5 shows how η depends on q_a for different values of the wall distance b . As the wall distance increases, not only does the mode amplitude grow, but also the unstable domain widens. The variation is strong for $b \leq 1.5$, but weak for $b > 1.5$. Thus we can approximately treat a




Figure 6: Lowest m components with $n = 1$ of the edge displacement obtained from free-boundary VMEC simulations with current profile $j(\rho) \propto (1 - \rho^8)$ and $\beta_N = 0.08\%$. $m = 4/n = 1$ is the dominant mode throughout the range of q_a , where the external kink mode is expected to be linearly unstable and non-linearly stable. The two black lines indicate the position of the marginal points computed with KINX.

wall distance of $b \gtrsim 1.5$ as if no wall was present. In the VMEC computations presented here, the field coils have a minimum distance of $b \approx 1.55$ to the plasma surface.

We find that the behaviour of η calculated from VMEC agrees well with the expected characteristics of external kink modes. In particular, the window of q_a over which η is non-zero, and also the shape




Figure 7: Amplitude of the saturated edge displacement of the 4/1 mode η_{41} computed from VMEC simulations at various values of β_N and comparison to the analytical model for current profile $j(\rho) \propto (1 - \rho^8)$.

of η with respect to q_a is roughly mirrored by the shape of γ with respect to q_a in Fig. 4a. Figure 6 shows the amplitude of η for each m (with $n = 1$), with η calculated according to Eq. (2). For an edge safety factor larger than the upper marginal point – given by the rational value $q_a = m/n$ – the plasma remains (nearly) axisymmetric. Lowering q_a below m/n results in 3D equilibria that have edge corrugations with external kink like properties as described previously. The amplitude of the saturated non-linear edge displacement is finite and dominated by a 4/1 component throughout the linearly unstable domain, as expected for the spectral properties of Fig. 3b. As q_a decreases below the lower marginal point, i.e. the value at which the linear external kink mode is stable, the displacement vanishes. For a small range of values of q_a slightly higher than the upper marginal point, or slightly lower than the lower marginal point we see small perturbations to η . We do not yet know the origin of these weak 3D saturated states.

A weak scaling of η with β_N is observed as shown in Fig. 7, where η_{41} calculated from VMEC at different values of β_N is compared with the analytically predicted non-linear saturated external kink amplitude η . The mode is present already at very low β_N , strongly indicating that the 3D states are dominantly current-driven, as expected for external kink modes. For the largest value of β_N the lower marginal point is shifted towards lower q_a , whereas it remains approximately constant for lower β_N values. Since finite beta effects are neglected in the analytical model, the most relevant comparison between VMEC and the non-linear analytic model is for small β_N in Fig. 7. Considering the shaped plasma and edge aspect ratio of




Figure 8: Mode spectrum (straight field line coordinates) of the VMEC edge displacement obtained in a simulation with $j(\rho) \propto (1 - \rho^4)^{1.1}$ and $q_a = 2.913$ at low pressure ($\beta_N = 0.08\%$). The $m = 3/n = 1$ mode is dominant.




Figure 9: Visualisation of the 3D equilibrium with current profile $j(\rho) \propto (1 - \rho^4)^{1.1}$, $q_a = 2.913$ and $\beta_N = 0.08\%$ shown at $\phi = \pi/3$. (a) Flux surfaces. (b) Perturbation $\delta p = p_{3D} - p_{2D}$ arising from the comparison of the pressure in the 3D equilibrium with the neighbouring state.

the VMEC JET-like simulations, the comparison with the cylindrical analytic model in Fig. 7 is very good.

Now that agreement between the VMEC displacement amplitude and the non-linear analytical external kink mode amplitude has been demonstrated, we further verify the results for a case with $q_a \lesssim 3$, for which a $n = 1, m = 3$ external mode is dominant. For this, in order to prevent the amplitude of the mode from becoming too large, we choose a less steep current profile $j(\rho) \propto (1 - \rho^4)^{1.1}$. As seen in Fig. 8, the 3/1 component (evaluated in sfl coordinates) is dominant in the edge mode spectrum of the resulting VMEC 3D equilibria as expected for a current-driven external kink mode. Figure 9 shows the flux surfaces and the 3D pressure perturbation, which are both non-axisymmetric and consistent with the mode spectrum. In Fig. 10 the analytically predicted saturated external kink amplitude is compared with the 3/1 component of the VMEC edge displacement at different values of




Figure 10: Non-linear amplitude of the 3/1 mode η_{31} obtained from VMEC with current profile $j(\rho) \propto (1 - \rho^4)^{1.1}$ and q_a around 3 at two values of β_N and comparison with the analytical prediction.

β_N . Again a weak scaling of η with the pressure is observed. The upper marginal point agrees well with the analytical value, whereas the lower one differs slightly. The saturated external kink amplitude peaks at a value of q_a closer to the lower marginal point, whereas the peak in the VMEC computations is somewhat shifted towards larger values. The reason for this is not currently certain, but it could be a result of finite aspect ratio and cross section shaping effects which are not taken into account in the analytic model.

4. Conclusions

Employing three-dimensional, free-boundary equilibrium computations, is shown to provide a novel way of obtaining saturated current-driven external kink modes. In principle, such saturated external states are time-independent and thus satisfy the force balance equation solved by an equilibrium code. For the first time, current-driven external kink modes are observed with an equilibrium code in free-boundary plasma configurations with standard monotonic current and q profiles typical of standard (baseline) tokamak plasma scenarios. Windows of the edge safety factor where external kink modes would be linearly unstable have been identified. In VMEC simulations with current profiles that result in an edge q value lying inside these regions, non-axisymmetric edge corrugations are observed. Analytical calculations of non-linear external kink modes reveal linear instability but non-linear stability, i.e. saturated mode amplitudes for the studied plasma configurations. To compare the analytic external kink amplitudes with VMEC, and thus verify that VMEC corrugations are those of standard non-linear saturated external kinks, the spectra of the VMEC

fluctuations are converted to spectra of a straight field line coordinate system. The Fourier spectrum calculated in this coordinate system shows one dominant mode, thus indicating consistency between 3D VMEC equilibria and analytical external kink models. Even though the mode amplitude scales weakly with β_N , the external perturbations seen in VMEC are of considerable size already at low β_N , indicating a current driven mode. Finite pressure is shown to have a weakly destabilising effect. The edge displacement in VMEC is found to be comparable to the analytically calculated saturated external kink mode amplitude. Small differences are observed in the value of q_a where the amplitude reaches a maximum. Due to the lack of the stabilizing effect of a conducting wall the 3D equilibrium simulations overestimate the saturated amplitude of a real tokamak by about 25% (for a wall distance $b = 1.2a$), the difference having been quantified via the analytic approach. We conclude that VMEC free-boundary calculations capture the salient features of saturated external kink modes, thus enabling efficient prediction of non-linear instability amplitudes, and e.g. fast ion and impurity transport studies.

Acknowledgements

This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 under grant agreement number 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission. This work was supported in part by the Swiss National Science Foundation. The authors thank Dr. S. P. Hirshman for providing us with the VMEC equilibrium code. We furthermore thank H. Eriksson for providing us with a corrected form of Eq. (16).

References

- [1] Freidberg J P 2014 *Ideal MHD* (Cambridge University Press)
- [2] Wesson J 1978 *Nuclear Fusion* **18** 87 URL <http://stacks.iop.org/0029-5515/18/i=1/a=010>
- [3] Eriksson H G and Wahlberg C 1994 *Plasma Physics and Controlled Fusion* **36** 1269 URL <http://stacks.iop.org/0741-3335/36/i=8/a=004>
- [4] Eriksson H G and Wahlberg C 1997 *Plasma Physics and Controlled Fusion* **39** 943
- [5] Brunetti D, Graves J, Cooper W and Terranova D 2014 *Nuclear Fusion* **54** 064017 URL <http://stacks.iop.org/0029-5515/54/i=6/a=064017>
- [6] Lütjens H and Luciani J F 2010 *Journal of Computational Physics* **229** 8130–8143
- [7] Marx A and Lütjens H 2017 *Computer Physics Communications* **212** 90 – 99 ISSN 0010-4655 URL <http://www.sciencedirect.com/science/article/pii/S0010465516303332>
- [8] Cooper W A, Graves J P, Duval B P, Porte L, Reimerdes H, Sauter O and Tran T M 2015 *Journal of Plasma*

- Physics* **81** 515810605 URL <https://doi.org/10.1017/S0022377815001221>
- [9] Cooper W A, Brunetti D, Duval B P, Faustin J M, Graves J P, Kleiner A, Patten H, Pfefferlé D, Porte L, Raghunathan M, Reimerdes H, Sauter O and Tran T M 2016 *Physics of Plasmas* **23** 040701 (Preprint <https://doi.org/10.1063/1.4945743>) URL <https://doi.org/10.1063/1.4945743>
- [10] Cooper W A, Graves J P, Duval B P, Sauter O, Faustin J M, Kleiner A, Lanthaler S, Patten H, Raghunathan M, Tran T M, Chapman I T and Ham C J 2016 *Plasma Physics and Controlled Fusion* **58** 064002 URL <http://stacks.iop.org/0741-3335/58/i=6/a=064002>
- [11] Pfefferlé D, Graves J, Cooper W, Misev C, Chapman I, Turnyanskiy M and Sangaroon S 2014 *Nuclear Fusion* **54** 064020 URL <http://stacks.iop.org/0029-5515/54/i=6/a=064020>
- [12] Raghunathan M, Graves J P, Nicolas T, Cooper W A, Garbet X and Pfefferlé D 2017 *Plasma Physics and Controlled Fusion* **59** 124002 URL <http://stacks.iop.org/0741-3335/59/i=12/a=124002>
- [13] Pfefferlé D, Cooper W, Graves J and Misev C 2014 *Computer Physics Communications* **185** 3127 – 3140 ISSN 0010-4655 URL <http://www.sciencedirect.com/science/article/pii/S0010465514002744>
- [14] Hirshman S P and Whitson J C 1983 *The Physics of Fluids* **26** 3553–3568 URL <http://aip.scitation.org/doi/abs/10.1063/1.864116>
- [15] Hirshman S, van RIJ W and Merkel P 1986 *Computer Physics Communications* **43** 143 – 155 ISSN 0010-4655 URL <http://www.sciencedirect.com/science/article/pii/0010465586900585>
- [16] Hanson J D and Hirshman S P 2002 *Physics of Plasmas* **9** 4410–4412 (Preprint <https://doi.org/10.1063/1.1507589>) URL <https://doi.org/10.1063/1.1507589>
- [17] Strumberger E, Günter S, Merkel P, Schwarz E and Tichmann C 2010 *Nuclear Fusion* **50** 025008 URL <http://stacks.iop.org/0029-5515/50/i=2/a=025008>
- [18] Degtyarev L, Martynov A, Medvedev S, Troyon F, Villard L and Gruber R 1997 *Computer Physics Communications* **103** 10 – 27 ISSN 0010-4655 URL <http://www.sciencedirect.com/science/article/pii/S0010465597000374>
- [19] Pao Y 1978 *The Physics of Fluids* **21** 765–772 (Preprint <http://aip.scitation.org/doi/pdf/10.1063/1.862296>) URL <http://aip.scitation.org/doi/abs/10.1063/1.862296>
- [20] Nakajima N 1990 *Physics of Fluids B: Plasma Physics* **2** 1170–1183 (Preprint <https://doi.org/10.1063/1.859253>) URL <https://doi.org/10.1063/1.859253>